

AUGMENTIN- amoxicillin and clavulanate potassium for suspension

Allegis Holdings, LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use AUGMENTIN safely and effectively. See full prescribing information for AUGMENTIN.

Augmentin (amoxicillin and clavulanate potassium) for oral suspension

Initial U.S. Approval: 1984

----- RECENT MAJOR CHANGES -----

Warnings and Precautions:

Drug-Induced Enterocolitis Syndrome (DIES) (5.3)

5/2024

----- INDICATIONS AND USAGE -----

AUGMENTIN is a combination of amoxicillin, a penicillin-class antibacterial and clavulanate potassium, a beta-lactamase inhibitor indicated for treatment of the following infections in adults and pediatric patients:

(1)

- Lower respiratory tract infections
- Acute bacterial otitis media
- Sinusitis
- Skin and skin structure infections
- Urinary tract infections

Limitations of Use

When susceptibility test results show susceptibility to amoxicillin, indicating no beta-lactamase production, AUGMENTIN should not be used. (1)

Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of AUGMENTIN and other antibacterial drugs, AUGMENTIN should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria. (1)

----- DOSAGE AND ADMINISTRATION -----

- Adults and Pediatric Patients greater than 40 kg: 500 or 875 mg every 12 hours or 250 or 500 mg every 8 hours, based on the amoxicillin component. (2.2, 2.3)
- Pediatric patients aged 12 weeks (3 months) and older: 25 to 45 mg/kg/day every 12 hours or 20 to 40 mg/kg/day every 8 hours, up to the adult dose. (2.3)
- Neonates and infants less than 12 weeks of age: 30 mg/kg/day divided every 12 hours, based on the amoxicillin component. Use of the 125 mg/5 mL oral suspension is recommended. (2.3)

----- DOSAGE FORMS AND STRENGTHS -----

- For Oral Suspension: 125 mg/31.25 mg per 5 mL, 250 mg/62.5 mg per 5 mL (3)

----- CONTRAINDICATIONS -----

- History of a serious hypersensitivity reaction (e.g., anaphylaxis or Stevens-Johnson syndrome) to AUGMENTIN or to other beta-lactams (e.g., penicillins or cephalosporins). (4.1)
- History of cholestatic jaundice/hepatic dysfunction associated with AUGMENTIN. (4.2)

----- WARNINGS AND PRECAUTIONS -----

- Serious (including fatal) hypersensitivity reactions: Discontinue AUGMENTIN if a reaction occurs. (5.1)
- Severe Cutaneous Adverse Reactions (SCAR): Monitor closely. Discontinue if rash progresses. (5.2)
- Drug-induced enterocolitis syndrome (DIES) has been reported with use of Amoxicillin, A component of Augmentin. If this occurs, discontinue AUGMENTIN and institute appropriate therapy (5.3)
- Hepatic dysfunction and cholestatic jaundice: Discontinue if signs/symptoms of hepatitis occur. Monitor liver function tests in patients with hepatic impairment. (5.4)
- *Clostridioides difficile*-associated diarrhea (CDAD): Evaluate patients if diarrhea occurs. (5.5)
- Patients with mononucleosis who receive AUGMENTIN develop skin rash. Avoid AUGMENTIN use in these patients. (5.6)
- Overgrowth: The possibility of superinfections with fungal or bacterial pathogens should be considered

during therapy. (5.7)

ADVERSE REACTIONS

5.4 The most frequently reported adverse effects were diarrhea/loose stools (9%), nausea (3%), skin rashes and urticaria (3%), vomiting (1%) and vaginitis (1%). (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Allegis Holdings, LLC at 1-866-633-9033 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Co-administration with probenecid is not recommended. (7.1)
- Concomitant use of AUGMENTIN and oral anticoagulants may increase the prolongation of prothrombin time.(7.2)
- Co-administration with allopurinol increases the risk of rash. (7.3)
- AUGMENTIN may reduce efficacy of oral contraceptives. (7.4)

USE IN SPECIFIC POPULATIONS

- Pediatric Use: Modify dose in patients 12 weeks or younger. (8.4)
- Renal Impairment: Dosage adjustment is recommended for severe renal impairment (GFR less than 30mL/min). (2.4, 8.6)]

Revised: 5/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

2.2 Adult Patients

2.3 Pediatric Patients

2.4 Patients with Renal Impairment

2.5 Directions for Mixing Amoxicillin and Clavulanate Potassium for Oral Suspension

2.6 Switching between Dosage Forms and between Strengths

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

4.1 Serious Hypersensitivity Reactions

4.2 Cholestatic Jaundice/Hepatic Dysfunction

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

5.2 Severe Cutaneous Adverse Reactions

5.3 Drug-Induced Enterocolitis Syndrome (DIES)

5.4 Hepatic Dysfunction

5.5 *Clostridioides difficile* Associated Diarrhea (CDAD)

5.6 Skin Rash in Patients with Mononucleosis

5.7 Potential for Microbial Overgrowth

5.8 Phenylketonurics

5.9 Development of Drug-Resistant Bacteria

6 ADVERSE REACTIONS

6.1 Clinical Trial Experience

6.2 Postmarketing Experience

7 DRUG INTERACTIONS

7.1 Probenecid

7.2 Oral Anticoagulants

- 7.3 Allopurinol
- 7.4 Oral Contraceptives
- 7.5 Effects on Laboratory Tests

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Labor and Delivery
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics
- 12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Lower Respiratory Tract and Complicated Urinary Tract Infections
- 14.2 Acute Bacterial Otitis Media and Diarrhea in Pediatric Patients

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

AUGMENTIN is indicated for the treatment of infections in adults and pediatric patients, due to susceptible isolates of the designated bacteria in the conditions listed below:

- **Lower Respiratory Tract Infections** - caused by beta-lactamase-producing isolates of *Haemophilus influenzae* and *Moraxella catarrhalis*.
- **Acute Bacterial Otitis Media** - caused by beta-lactamase-producing isolates of *H. influenzae* and *M. catarrhalis*.
- **Sinusitis** - caused by beta-lactamase-producing isolates of *H. influenzae* and *M. catarrhalis*.
- **Skin and Skin Structure Infections** - caused by beta-lactamase-producing isolates of *Staphylococcus aureus*, *Escherichia coli*, and *Klebsiella* species.
- **Urinary Tract Infections** - caused by beta-lactamase-producing isolates of *E. coli*, *Klebsiella* species, and *Enterobacter* species.

Limitations of Use

When susceptibility test results show susceptibility to amoxicillin, indicating no beta-lactamase production, AUGMENTIN should not be used.

Usage

To reduce the development of drug-resistant bacteria and maintain the effectiveness of AUGMENTIN and other antibacterial drugs, AUGMENTIN should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

2 DOSAGE AND ADMINISTRATION

2.1 Important Administration Instructions

AUGMENTIN may be taken without regard to meals; however, absorption of clavulanate potassium is enhanced when AUGMENTIN is administered at the start of a meal. To minimize the potential for gastrointestinal intolerance, AUGMENTIN should be taken at the start of a meal.

2.2 Adult Patients

See dosing regimens of AUGMENTIN (based on the amoxicillin component) provided in Table 1 below.

Table 1. Dosing Regimens of AUGMENTIN in Adult Patients

<u>TYPE OF INFECTION</u>	<u>DOSING REGIMEN OF Amoxicillin and Clavulanate Potassium</u>
Severe infections and infections of the respiratory tract	one 875 mg tablet ^a of AUGMENTIN every 12 hours or one 500 mg tablet ^{b,c} of AUGMENTIN every 8 hours
Less severe infections	one 500 mg tablet ^{b,c} of AUGMENTIN every 12 hours or one 250 mg tablet ^d of AUGMENTIN every 8 hours

^a Adults who have difficulty swallowing may be given the AUGMENTIN 200 mg/28.5 mg per 5 mL suspension or the AUGMENTIN 400 mg/57 mg per 5 mL suspension may be used in place of the 875 mg/125 mg tablet.

^b Adults who have difficulty swallowing may be given the AUGMENTIN 125 mg/31.25 mg per 5 mL or AUGMENTIN 250 mg/62.5 mg per 5 mL suspension in place of the 500 mg/125 mg tablet.

^c Two AUGMENTIN 250 mg/125 mg tablets are *NOT* substitutable with one 500 mg/125 mg AUGMENTIN tablet [see *Dosage and Administration* (2.6)] .

^d AUGMENTIN 250 mg/125 mg tablet is *NOT* substitutable with AUGMENTIN 250 mg/62.5 mg chewable tablet [see *Dosage and Administration* (2.6)]

2.3 Pediatric Patients

Based on the amoxicillin component, AUGMENTIN should be dosed as follows:

Neonates and Infants Aged less than 12 weeks (less than 3 months): See dosing regimens of AUGMENTIN provided in Table 2 below.

Table 2: Dosing Regimens of AUGMENTIN in Neonates and Infants Aged Less than 12 Weeks (Less than 3 Months)

<u>PATIENT POPULATION</u>	<u>DOSING REGIMEN</u>
	<u>AUGMENTIN 125 mg/31.25 mg per 5 mL for oral suspension^a</u>
Neonates and Infants aged less than 12 weeks (less than 3 months)	30 mg/kg/day every 12 hours

^a Experience with the AUGMENTIN 200 mg/28.5 mg per 5 mL formulation in this age group is limited, and thus, use of the AUGMENTIN 125 mg/31.25 mg per 5 mL for oral suspension is recommended.

Patients Aged 12 weeks (3 months) and Older and Weighing Less than 40 kg: See dosing regimens provided in Table 3 below.

The every 12 hour regimen is recommended as it is associated with significantly less diarrhea [see *Clinical Studies* (14.2)] .

The Amoxicillin and Clavulanate Potassium 200 mg/28.5 mg per 5 mL and Amoxicillin and Clavulanate Potassium 400 mg/57 mg per 5 mL) for oral suspension and Amoxicillin and Clavulanate Potassium 200 mg/28.5 mg and Amoxicillin and Clavulanate Potassium 400 mg/57 mg chewable tablets contain aspartame and should not be used by phenylketonurics. [see *Warnings and Precautions* (5.8)].

Table 3: Dosing in Patients Aged 12 Weeks (3 Months) and Older and Weighing Less than 40 kg

INFECTION	DOSING REGIMEN	
	Every 12 hours	Every 8 hours
	Amoxicillin and Clavulanate Potassium 200 mg/28.5 mg per 5 mL or Amoxicillin and Clavulanate Potassium 400 mg/57 mg per 5 mL oral suspension ^a	Amoxicillin and Clavulanate Potassium 125 mg/31.25 mg per 5 mL or Amoxicillin and Clavulanate Potassium 250 mg/62.5 mg per 5 mL oral suspension ^a
Otitis media ^b , sinusitis, lower respiratory tract infections, and more severe infections	45 mg/kg/day every 12 hours	40 mg/kg/day every 8 hours
Less severe infections	25 mg/kg/day every 12 hours	20 mg/kg/day every 8 hours

^a Each strength of Amoxicillin and Clavulanate Potassium oral suspension is available as a chewable tablet for use by older children.

^b Duration of therapy studied and recommended for acute otitis media is 10 days.

Patients Weighing 40 kg or More: Pediatric patients weighing 40 kg or more should be dosed according to adult recommendations.

- The 250 mg/125 mg tablet of AUGMENTIN should *NOT* be used until the child weighs at least 40 kg, due to the different amoxicillin to clavulanic acid ratios in the 250 mg/125 mg tablet of AUGMENTIN versus the 250 mg/62.5 mg chewable tablet of AUGMENTIN.

2.4 Patients with Renal Impairment

Patients with impaired renal function do not generally require a reduction in dose unless the impairment is severe. Renal impairment patients with a glomerular filtration rate (GFR) of less than 30 mL/min should *NOT* receive the 875 mg dose (based on the amoxicillin component) of AUGMENTIN. See dosing regimens in patients with severe renal impairment provided in Table 4.

Table 4. Dosing Regimens of AUGMENTIN in Patients with Severe Renal Impairment

Patients with Renal Impairment	Dosing Regimen
GFR 10 mL/min to 30 mL/min	500 mg or 250 mg every 12 hours, depending on the severity of the infection
GFR less than 10 mL/min	500 mg or 250 mg every 24 hours, depending on severity of the infection
Hemodialysis	500 mg or 250 mg every 24 hours, depending on severity of the infection Administer an additional dose both during and at the end of dialysis

2.5 Directions for Mixing Amoxicillin and Clavulanate Potassium for Oral Suspension

Prepare AUGMENTIN for oral suspension at time of dispensing as follows: Tap bottle until all powder flows freely. Measure a total (see Table 5 below for total amount of water for reconstitution) OF WATER. Add approximately 2/3 of the water to the powder. Replace cap and shake VIGOROUSLY. Add remaining water. Replace cap and shake VIGOROUSLY.

Table 5: Amount of Water for Mixing AUGMENTIN for Oral Suspension

Strength of AUGMENTIN Oral Suspension	Bottle Size	Amount of Water for Reconstitution	Contents of Each Teaspoonful (5 mL)
125 mg/31.25 mg per 5 mL	75 mL	67 mL	125 mg of amoxicillin and 31.25 mg of clavulanic acid as the potassium

	100 mL	90 mL	salt
	150 mL	134 mL	
250 mg/62.5 mg per 5 mL	75 mL	65 mL	250 mg of amoxicillin and 62.5 mg of clavulanic acid as the potassium salt
	100 mL	87 mL	
	150 mL	130 mL	

Shake oral suspension well before using. Reconstituted suspension must be stored under refrigeration and discarded after 10 days. Some color change is normal during dosing period.

2.6 Switching between Dosage Forms and between Strengths

AUGMENTIN 250 mg/125 mg Tablet is NOT Substitutable with AUGMENTIN 250 mg/62.5 mg Chewable Tablet

The 250 mg/125 mg tablet of AUGMENTIN and the 250 mg/62.5 mg chewable tablet of AUGMENTIN should *NOT* be substituted for each other and the 250 mg/125 mg tablet of AUGMENTIN should *NOT* be used in pediatric patients weighing less than 40 kg [see *Dosage and Administration (2.3)*]. The 250 mg tablet of AUGMENTIN and the 250 mg chewable tablet of AUGMENTIN do not contain the same amount of clavulanic acid. The 250 mg tablet of AUGMENTIN contains 125 mg of clavulanic acid whereas the 250 mg chewable tablet of AUGMENTIN contains 62.5 mg of clavulanic acid.

Two AUGMENTIN 250 mg/125 mg Tablets are NOT Substitutable with One 500 mg/125 mg AUGMENTIN Tablet

Two 250 mg/125 mg tablets of AUGMENTIN should *NOT* be substituted for one 500 mg/125 mg tablet of AUGMENTIN. Since both the 250 mg and 500 mg tablets of AUGMENTIN contain the same amount of clavulanic acid (125 mg, as the potassium salt), two 250 mg tablets of AUGMENTIN are not equivalent to one 500 mg tablet of AUGMENTIN.

3 DOSAGE FORMS AND STRENGTHS

Augmentin for Oral Suspension, USP:

- **125 mg/31.25 mg per 5 mL:** Banana-flavored powder for oral suspension (each 5 mL of reconstituted suspension contains 125 mg of amoxicillin as the trihydrate and 31.25 mg of clavulanic acid as the potassium salt).
- **250 mg/62.5 mg per 5 mL:** Orange-flavored powder for oral suspension (each 5 mL of reconstituted suspension contains 250 mg of amoxicillin as the trihydrate and 62.5 mg of clavulanic acid as the potassium salt).

4 CONTRAINDICATIONS

4.1 Serious Hypersensitivity Reactions

AUGMENTIN is contraindicated in patients with a history of serious hypersensitivity reactions (e.g., anaphylaxis or Stevens-Johnson syndrome) to amoxicillin, clavulanate or to other beta-lactam antibacterial drugs (e.g., penicillins and cephalosporins).

4.2 Cholestatic Jaundice/Hepatic Dysfunction

AUGMENTIN is contraindicated in patients with a previous history of cholestatic jaundice/hepatic dysfunction associated with AUGMENTIN.

5 WARNINGS AND PRECAUTIONS

5.1 Hypersensitivity Reactions

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions have been reported in patients receiving beta-lactam antibacterials, including AUGMENTIN. These reactions are more likely to occur in individuals with a history of penicillin hypersensitivity and/or a history of sensitivity to multiple allergens. Before initiating therapy with AUGMENTIN, careful inquiry should be made regarding previous hypersensitivity reactions to penicillins, cephalosporins, or other allergens. If an allergic reaction occurs, AUGMENTIN should be discontinued, and appropriate therapy instituted.

5.2 Severe Cutaneous Adverse Reactions

AUGMENTIN may cause severe cutaneous adverse reactions (SCAR), such as Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS), and acute generalized exanthematous pustulosis (AGEP). If patients develop a skin rash, they should be monitored closely, and AUGMENTIN discontinued if lesions progress.

5.3 Drug-Induced Enterocolitis Syndrome (DIES)

Drug-induced enterocolitis syndrome (DIES) has been reported with use of amoxicillin, a component of AUGMENTIN

[see Adverse Reactions (6.2)], with most cases occurring in pediatric patients ≤ 18 years of age. DIES is a non IgE mediated

hypersensitivity reaction characterized by protracted vomiting occurring 1 to 4 hours after drug ingestion in the absence of

skin or respiratory symptoms. DIES may be associated with pallor, lethargy, hypotension, shock, diarrhea within 24 hours of ingesting

amoxicillin, and leukocytosis with neutrophilia. If DIES occurs, discontinue AUGMENTIN and institute appropriate therapy.

5.4 Hepatic Dysfunction

Hepatic dysfunction, including hepatitis and cholestatic jaundice has been associated with the use of AUGMENTIN. Hepatic toxicity is usually reversible; however, deaths have been reported. Hepatic function should be monitored at regular intervals in patients with

hepatic impairment.

5.5 *Clostridioides difficile* Associated Diarrhea (CDAD)

Clostridioides difficile associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents, including AUGMENTIN, and may range in severity from mild diarrhea to fatal colitis. Treatment with antibacterial agents alters the normal flora of the colon leading to overgrowth of *C. difficile*.

C. difficile produces toxins A and B which contribute to the development of CDAD. Hypertoxin-producing strains of *C. difficile* cause increased morbidity and mortality, as these infections can be refractory to antimicrobial therapy and may require colectomy. CDAD must be considered in all patients who present with diarrhea following antibacterial use. Careful medical history is necessary since CDAD has been reported to occur over 2 months after the administration of antibacterial agents.

If CDAD is suspected or confirmed, ongoing antibacterial use not directed against *C. difficile* may need to be discontinued. Appropriate fluid and electrolyte management, protein supplementation, antibacterial treatment of *C. difficile*, and surgical evaluation should be instituted as clinically indicated.

5.6 Skin Rash in Patients with Mononucleosis

A high percentage of patients with mononucleosis who receive amoxicillin develop an erythematous skin rash. Thus, AUGMENTIN should not be administered to patients with mononucleosis.

5.7 Potential for Microbial Overgrowth

The possibility of superinfections with fungal or bacterial pathogens should be considered during therapy. If superinfection occurs, amoxicillin and clavulanate potassium should be discontinued and appropriate therapy instituted.

5.8 Phenylketonurics

AUGMENTIN Chewable tablets and AUGMENTIN for Oral Suspension contain aspartame which contains phenylalanine. Each 200 mg chewable tablet of AUGMENTIN contains 2.1 mg phenylalanine; each 400 mg chewable tablet contains 4.2 mg phenylalanine; each 5 mL of either the 200 mg/5 mL or 400 mg/5 mL oral suspension contains 7 mg phenylalanine. The other formulations of AUGMENTIN do not contain phenylalanine.

5.9 Development of Drug-Resistant Bacteria

Prescribing AUGMENTIN in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

6 ADVERSE REACTIONS

The following are discussed in more detail in other sections of the labeling:

- Anaphylactic reactions [see *Warnings and Precautions (5.1)*]
- Severe Cutaneous Adverse Reactions [see *Warnings and Precautions (5.2)*]
- Drug-Induced Enterocolitis Syndrom (DIES) [see *Warnings and Precautions (5.3)*]

- Hepatic Dysfunction [see *Warnings and Precautions* (5.4)]
- *Clostridioides difficile* Associated Diarrhea (CDAD) [see *Warnings and Precautions* (5.5)]

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The most frequently reported adverse reactions were diarrhea/loose stools (9%), nausea (3%), skin rashes and urticaria (3%), vomiting (1%) and vaginitis (1%). Less than 3% of patients discontinued therapy because of drug-related adverse reactions. The overall incidence of adverse reactions, and in particular diarrhea, increased with the higher recommended dose. Other less frequently reported adverse reactions (less than 1%) include: Abdominal discomfort, flatulence, and headache.

In pediatric patients (aged 2 months to 12 years), 1 US/Canadian clinical trial was conducted which compared 45/6.4 mg/kg/day (divided every 12 hours) of Amoxicillin and Clavulanate Potassium for 10 days versus 40/10 mg/kg/day (divided every 8 hours) of Amoxicillin and Clavulanate Potassium for 10 days in the treatment of acute otitis media. A total of 575 patients were enrolled, and only the suspension formulations were used in this trial. Overall, the adverse reactions seen were comparable to that noted above; however, there were differences in the rates of diarrhea, skin rashes/urticaria, and diaper area rashes. [see *Clinical Studies* (14.2)] .

6.2 Postmarketing Experience

In addition to adverse reactions reported from clinical trials, the following have been identified during postmarketing use of AUGMENTIN. Because they are reported voluntarily from a population of unknown size, estimates of frequency cannot be made. These events have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to Amoxicillin and Clavulanate Potassium.

Gastrointestinal: Drug-induced enterocolitis syndrome (DIES), indigestion, gastritis, stomatitis, glossitis, black “hairy” tongue, mucocutaneous candidiasis, enterocolitis, and hemorrhagic/pseudomembranous colitis. Onset of pseudomembranous colitis symptoms may occur during or after antibacterial treatment. [see *Warnings and Precautions* (5.5)] .

Immune: Hypersensitivity reactions, anaphylactic/anaphylactoid reactions (including shock), angioedema, serum sickness-like reactions (urticaria or skin rash accompanied by arthritis, arthralgia, myalgia, and frequently fever), hypersensitivity vasculitis [see *Warnings and Precautions* (5.1)] .

Skin and Appendages: Rashes, pruritus, urticaria, erythema multiforme, SJS, TEN, DRESS, AGEP, exfoliative dermatitis , and *linear IgA dermatosis*.

Liver: Hepatic dysfunction, including hepatitis and cholestatic jaundice, increases in serum transaminases (AST and/or ALT), serum bilirubin, and/or alkaline phosphatase, has been reported with Amoxicillin and Clavulanate Potassium. It has been reported more commonly in the elderly, in males, or in patients on prolonged treatment. The histologic findings on liver biopsy have consisted of predominantly cholestatic,

hepatocellular, or mixed cholestatic hepatocellular changes. The onset of signs/symptoms of hepatic dysfunction may occur during or several weeks after therapy has been discontinued. The hepatic dysfunction, which may be severe, is usually reversible. Deaths have been reported [see *Contraindications (4.2)*, *Warnings and Precautions (5.4)*].

Renal: Interstitial nephritis, hematuria, and crystalluria have been reported [see *Overdosage (10)*].

Hemic and Lymphatic Systems: Anemia, including hemolytic anemia, thrombocytopenia, thrombocytopenic purpura, eosinophilia, leukopenia, and agranulocytosis have been reported. These reactions are usually reversible on discontinuation of therapy and are believed to be hypersensitivity phenomena. Thrombocytosis was noted in less than 1% of the patients treated with AUGMENTIN. There have been reports of increased prothrombin time in patients receiving AUGMENTIN and anticoagulant therapy concomitantly [see *Drug Interactions (7.2)*].

Central Nervous System: Agitation, anxiety, behavioral changes, aseptic meningitis, confusion, convulsions, dizziness, insomnia, and reversible hyperactivity have been reported.

Miscellaneous: Tooth discoloration (brown, yellow, or gray staining) has been reported. Most reports occurred in pediatric patients. Discoloration was reduced or eliminated with brushing or dental cleaning in most cases.

7 DRUG INTERACTIONS

7.1 Probenecid

Probenecid decreases the renal tubular secretion of amoxicillin but does not delay renal excretion of clavulanic acid. Concurrent use with AUGMENTIN may result in increased and prolonged blood concentrations of amoxicillin. Co-administration of probenecid is not recommended.

7.2 Oral Anticoagulants

Abnormal prolongation of prothrombin time (increased international normalized ratio [INR]) has been reported in patients receiving amoxicillin and oral anticoagulants. Appropriate monitoring should be undertaken when anticoagulants are prescribed concurrently with AUGMENTIN. Adjustments in the dose of oral anticoagulants may be necessary to maintain the desired level of anticoagulation.

7.3 Allopurinol

The concurrent administration of allopurinol and amoxicillin increases the incidence of rashes in patients receiving both drugs as compared to patients receiving amoxicillin alone. It is not known whether this potentiation of amoxicillin rashes is due to allopurinol or the hyperuricemia present in these patients.

7.4 Oral Contraceptives

AUGMENTIN may affect intestinal flora, leading to lower estrogen reabsorption and reduced efficacy of combined oral estrogen/progesterone contraceptives.

7.5 Effects on Laboratory Tests

High urine concentrations of amoxicillin may result in false-positive reactions when testing for the presence of glucose in urine using CLINITEST[®], Benedict's Solution, or Fehling's Solution. Since this effect may also occur with AUGMENTIN, it is recommended that glucose tests based on enzymatic glucose oxidase reactions be used.

Following administration of amoxicillin to pregnant women, a transient decrease in plasma concentration of total conjugated estriol, estriol-glucuronide, conjugated estrone, and estradiol has been noted.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Teratogenic Effects: Pregnancy Category B. Reproduction studies performed in pregnant rats and mice given AUGMENTIN (2:1 ratio formulation of amoxicillin:clavulanate) at oral doses up to 1200 mg/kg/day revealed no evidence of harm to the fetus due to AUGMENTIN. The amoxicillin doses in rats and mice (based on body surface area) were approximately 4 and 2 times the maximum recommended adult human oral dose (875 mg every 12 hours). For clavulanate, these dose multiples were approximately 9 and 4 times the maximum recommended adult human oral dose (125 mg every 8 hours). There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

8.2 Labor and Delivery

Oral ampicillin-class antibacterials are poorly absorbed during labor. It is not known whether use of amoxicillin and clavulanate potassium in humans during labor or delivery has immediate or delayed adverse effects on the fetus, prolongs the duration of labor, or increases the likelihood of the necessity for an obstetrical intervention.

8.3 Nursing Mothers

Amoxicillin has been shown to be excreted in human milk. Amoxicillin and clavulanate potassium use by nursing mothers may lead to sensitization of infants. Caution should be exercised when amoxicillin and clavulanate potassium is administered to a nursing woman.

8.4 Pediatric Use

The safety and effectiveness of AUGMENTIN for Oral Suspension and Chewable Tablets have been established in pediatric patients. Use of AUGMENTIN in pediatric patients is supported by evidence from studies of AUGMENTIN Tablets in adults with additional data from a study of AUGMENTIN for Oral Suspension in pediatric patients aged 2 months to 12 years with acute otitis media [see *Clinical Studies* (14.2)] .

Because of incompletely developed renal function in neonates and young infants, the elimination of amoxicillin may be delayed; clavulanate elimination is unaltered in this age group. Dosing of Amoxicillin and Clavulanate Potassium should be modified in pediatric patients aged less than 12 weeks (less than 3 months) [see *Dosage and Administration* (

2.3)].

8.5 Geriatric Use

Of the 3,119 patients in an analysis of clinical studies of AUGMENTIN, 32% were greater than or equal to 65 years old, and 14% were greater than or equal to 75 years old. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

This drug is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

8.6 Renal Impairment

Amoxicillin is primarily eliminated by the kidney and dosage adjustment is usually required in patients with severe renal impairment (GFR less than 30 mL/min). See Patients with Renal Impairment [see *Dosage and Administration* (2.4)] for specific recommendations in patients with renal impairment.

10 OVERDOSAGE

In case of overdosage, discontinue medication, treat symptomatically, and institute supportive measures as required. A prospective study of 51 pediatric patients at a poison-control center suggested that overdosages of less than 250 mg/kg of amoxicillin are not associated with significant clinical symptoms ¹.

Interstitial nephritis resulting in oliguric renal failure has been reported in patients after overdosage with amoxicillin and clavulanate potassium.

Crystalluria, in some cases leading to renal failure, has also been reported after amoxicillin and clavulanate potassium overdosage in adult and pediatric patients. In case of overdosage, adequate fluid intake and diuresis should be maintained to reduce the risk of amoxicillin and clavulanate potassium crystalluria.


Renal impairment appears to be reversible with cessation of drug administration. High blood levels may occur more readily in patients with impaired renal function because of decreased renal clearance of amoxicillin and clavulanate potassium. Amoxicillin and clavulanate potassium may be removed from circulation by hemodialysis [see *Dosage and Administration* (2.4)] .

11 DESCRIPTION


AUGMENTIN is an oral antibacterial combination consisting of amoxicillin and the beta-lactamase inhibitor, clavulanate potassium (the potassium salt of clavulanic acid).

Amoxicillin is an analog of ampicillin, derived from the basic penicillin nucleus, 6-aminopenicillanic acid. The amoxicillin molecular formula is $C_{16}H_{19}N_3O_5S \cdot 3H_2O$, and the molecular weight is 419.46. Chemically, amoxicillin is (2*S*,5*R*,6*R*)-6-[(*R*)-(-)-2-Amino-2-(*p*-hydroxyphenyl)acetamido]-3,3-dimethyl-7-oxo-4-thia-1-

azabicyclo[3.2.0]heptane-2-carboxylic acid trihydrate and may be represented structurally as:



Clavulanic acid is produced by the fermentation of *Streptomyces clavuligerus*. It is a beta-lactam structurally related to the penicillins and possesses the ability to inactivate some beta-lactamases by blocking the active sites of these enzymes. The clavulanate potassium molecular formula is $C_8H_8KNO_5$, and the molecular weight is 237.25. Chemically, clavulanate potassium is potassium (Z)(2R,5R)-3-(2-hydroxyethylidene)-7-oxo-4-oxa-1-azabicyclo[3.2.0]-heptane-2-carboxylate and may be represented structurally as:



AUGMENTIN for Oral Suspension:

- **125 mg/31.25 mg:** Following constitution, each 5 mL of oral suspension contains 125 mg of amoxicillin as the trihydrate, and 31.25 mg of clavulanic acid (equivalent to 37.23 mg of clavulanate potassium).
- **250 mg/62.5 mg:** Following constitution, each 5 mL of oral suspension contains 250 mg of amoxicillin as the trihydrate, and 62.5 mg of clavulanic acid (equivalent to 74.5 mg of clavulanate potassium).

Inactive Ingredients:

- **AUGMENTIN for Oral Suspension, 125 mg/31.5 mg per 5mL and 250 mg/62.5 mg per 5mL** - Colloidal silicon dioxide, flavorings, xanthan gum, mannitol, succinic acid, silica gel and sodium saccharin.
- Each 5 mL of reconstituted 125 mg/31.5 mg oral suspension of AUGMENTIN contains 0.16 mEq potassium
- Each 5 mL of reconstituted 250 mg/62.5 mg oral suspension of AUGMENTIN contains 0.32 mEq potassium

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

AUGMENTIN is an antibacterial drug [see *Microbiology* (12.4)] .

12.3 Pharmacokinetics

Mean amoxicillin and clavulanate potassium pharmacokinetic parameters in normal adults following administration of Amoxicillin and Clavulanate Potassium Tablets are shown in Table 6 and following administration of Amoxicillin and Clavulanate Potassium for Oral Suspension and Chewable Tablets are shown in Table 7.

Table 6: Mean (\pm S.D.) Amoxicillin and Clavulanate Potassium Pharmacokinetic Parameters a,b with Amoxicillin and Clavulanate Potassium Tablets

Dose and Regimen of Amoxicillin and Clavulanate	C _{max} (mcg/mL)		AUC ₀₋₂₄ (mcg*h/mL)	
	Amoxicillin	Clavulanate potassium	Amoxicillin	Clavulanate potassium
250 mg/125 mg every 8 hours	3.3 \pm 1.12	1.5 \pm 0.70	26.7 \pm 4.56	12.6 \pm 3.25
500 mg/125 mg every 12 hours	6.5 \pm 1.41	1.8 \pm 0.61	33.4 \pm 6.76	8.6 \pm 1.95
500 mg/125 mg every 8 hours	7.2 \pm 2.26	2.4 \pm 0.83	53.4 \pm 8.87	15.7 \pm 3.86
875 mg/125 mg every 12 hours	11.6 \pm 2.78	2.2 \pm 0.99	53.5 \pm 12.31	10.2 \pm 3.04

^a Mean (\pm standard deviation) values of 14 normal adults (N equals 15 for clavulanate potassium in the low-dose regimens). Peak concentrations occurred approximately 1.5 hours after the dose.

^b Amoxicillin and clavulanate potassium administered at the start of a light meal.

Table 7: Mean (\pm S.D.) Amoxicillin and Clavulanate Potassium Pharmacokinetic Parameters a,b with Amoxicillin and Clavulanate Potassium for Oral Suspension and Chewable Tablets

Dose of Amoxicillin and Clavulanate	C _{max} (mcg/mL)		AUC ₀₋₂₄ (mcg*h/mL)	
	Amoxicillin	Clavulanate potassium	Amoxicillin	Clavulanate potassium
400 mg/57 mg (5 mL of suspension)	6.94 \pm 1.24	1.10 \pm 0.42	17.29 \pm 2.28	2.34 \pm 0.94
400 mg/57 mg (1 chewable tablet)	6.67 \pm 1.37	1.03 \pm 0.33	17.24 \pm 2.64	2.17 \pm 0.73

^a Mean (\pm standard deviation) values of 28 normal adults. Peak concentrations occurred approximately 1 hour after the dose.

^b Amoxicillin and clavulanate potassium administered at the start of a light meal.

Oral administration of 5 mL of the 250 mg/62.5 mg suspension of Amoxicillin and Clavulanate Potassium or the equivalent dose of 10 mL of the 125 mg/31.25 mg suspension of Amoxicillin and Clavulanate Potassium provides average peak serum concentrations approximately 1 hour after dosing of 6.9 mcg/mL for amoxicillin and 1.6 mcg/mL for clavulanic acid. The areas under the serum concentration curves obtained during the first 4 hours after dosing were 12.6 mcg*h/mL for amoxicillin and 2.9 mcg*h/mL for clavulanic acid when 5 mL of the 250 mg/62.5 mg suspension of Amoxicillin and Clavulanate Potassium or equivalent dose of 10 mL of the 125 mg/31.25 mg suspension of Amoxicillin and Clavulanate Potassium were administered to normal adults. One 250 mg/62.5 mg chewable tablet of Amoxicillin and Clavulanate Potassium or two 125 mg/31.25 mg chewable tablets of Amoxicillin and Clavulanate Potassium are equivalent to 5 mL of the 250 mg/62.5 mg suspension of Amoxicillin and Clavulanate Potassium and provide similar serum concentrations of amoxicillin and clavulanic acid.

Amoxicillin serum concentrations achieved with Amoxicillin and Clavulanate Potassium are similar to those produced by the oral administration of equivalent doses of amoxicillin alone. Time above the minimum inhibitory concentration of 1 mcg/mL for amoxicillin has been shown to be similar after corresponding every 12 hour and every 8-hour dosing regimens of Amoxicillin and Clavulanate Potassium in adults and children.

Absorption: Dosing in the fasted or fed state has minimal effect on the pharmacokinetics of amoxicillin. While AUGMENTIN can be given without regard to meals, absorption of clavulanate potassium when taken with food is greater relative to the fasted state. In one study, the relative bioavailability of clavulanate was reduced when AUGMENTIN was dosed at 30 and 150 minutes after the start of a high-fat breakfast.

Distribution: Neither component in AUGMENTIN is highly protein-bound; clavulanic acid is approximately 25% bound to human serum and amoxicillin approximately 18% bound. Amoxicillin diffuses readily into most body tissues and fluids with the exception of the brain and spinal fluid.

Two hours after oral administration of a single 35 mg/kg dose of suspension of AUGMENTIN to fasting children, average concentrations of 3 mcg/mL of amoxicillin and 0.5 mcg/mL of clavulanic acid were detected in middle ear effusions.

Metabolism and Excretion: The half-life of amoxicillin after the oral administration of AUGMENTIN is 1.3 hours and that of clavulanic acid is 1 hour.

Approximately 50% to 70% of the amoxicillin and approximately 25% to 40% of the clavulanic acid are excreted unchanged in urine during the first 6 hours after administration of a single 250 mg/125 mg or 500 mg/125 mg tablet of AUGMENTIN.

12.4 Microbiology

Amoxicillin is a semisynthetic antibacterial with in vitro bactericidal activity against Gram-positive and Gram-negative bacteria. Amoxicillin is, however, susceptible to degradation by beta-lactamases, and therefore, the spectrum of activity does not include organisms which produce these enzymes. Clavulanic acid is a beta-lactam, structurally related to the penicillins, which possesses the ability to inactivate some beta-lactamase enzymes commonly found in microorganisms resistant to penicillins and cephalosporins. In particular, it has good activity against the clinically important plasmid-mediated beta-lactamases frequently responsible for transferred drug resistance.

The formulation of amoxicillin and clavulanic acid in AUGMENTIN protects amoxicillin from degradation by some beta-lactamase enzymes and extends the antibacterial spectrum of amoxicillin to include many bacteria normally resistant to amoxicillin.

Amoxicillin and clavulanic acid has been shown to be active against most isolates of the following bacteria, both in vitro and in clinical infections [*see Indications and Usage (1)*] .

Gram-positive bacteria

Staphylococcus aureus

Gram-negative bacteria

Enterobacter species

Escherichia coli

Haemophilus influenzae

Klebsiella species

Moraxella catarrhalis

The following in vitro data are available, **but their clinical significance is unknown**. At least 90 percent of the following bacteria exhibit an in vitro minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for amoxicillin and clavulanic acid. However, the efficacy of amoxicillin and clavulanic acid in treating clinical infections due to these bacteria **has not been** established in adequate and well-controlled clinical trials.

Gram-positive bacteria

Enterococcus faecalis

Staphylococcus epidermidis

Staphylococcus saprophyticus

Streptococcus pneumoniae

Streptococcus pyogenes

Viridans group *Streptococcus*

Gram-negative Bacteria

Eikenella corrodens

Proteus mirabilis

Anaerobic Bacteria

Bacteroides species including *Bacteroides fragilis*

Fusobacterium species

Peptostreptococcus species

Susceptibility Test Methods

For specific information regarding susceptibility test interpretive criteria and associated test methods and quality control standards recognized by FDA for this drug, please see: <https://www.fda.gov/STIC>.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies in animals have not been performed to evaluate carcinogenic potential.

AUGMENTIN (4:1 ratio formulation of amoxicillin:clavulanate) was non-mutagenic in the Ames bacterial mutation assay, and the yeast gene conversion assay. AUGMENTIN was weakly positive in the mouse lymphoma assay, but the trend toward increased mutation frequencies in this assay occurred at doses that were also associated with decreased cell survival. AUGMENTIN was negative in the mouse micronucleus test, and in the dominant lethal assay in mice. Potassium clavulanate alone was tested in the Ames bacterial mutation assay and in the mouse micronucleus test, and was negative in each

of these assays.

AUGMENTIN (2:1 ratio formulation of amoxicillin:clavulanate) at oral doses of up to 1,200 mg/kg/day was found to have no effect on fertility and reproductive performance in rats. Based on body surface area, this dose of amoxicillin is approximately 4 times the maximum recommended adult human oral dose (875 mg every 12 hours). For clavulanate, the dose multiple is approximately 9 times higher than the maximum recommended adult human oral dose (125 mg every 8 hours), also based on body surface area.

14 CLINICAL STUDIES

14.1 Lower Respiratory Tract and Complicated Urinary Tract Infections

Data from 2 pivotal trials in 1,191 patients treated for either lower respiratory tract infections or complicated urinary tract infections compared a regimen of 875 mg/125 mg tablets of Amoxicillin and Clavulanate Potassium every 12 hours to 500 mg/125 mg tablets of Amoxicillin and Clavulanate Potassium dosed every 8 hours (584 and 607 patients, respectively). Comparable efficacy was demonstrated between the every 12 hours and every 8 hours dosing regimens. There was no significant difference in the percentage of adverse events in each group. The most frequently reported adverse event was diarrhea; incidence rates were similar for the 875 mg/125 mg every 12 hours and 500 mg/125 mg every 8 hours dosing regimens (15% and 14%, respectively); however, there was a statistically significant difference (p less than 0.05) in rates of severe diarrhea or withdrawals with diarrhea between the regimens: 1% for 875 mg/125 mg every 12 hours regimen versus 2% for the 500 mg/125 mg every 8 hours regimen.

In one of these pivotal trials, patients with either pyelonephritis (n equals 361) or a complicated urinary tract infection (i.e., patients with abnormalities of the urinary tract that predispose to relapse of bacteriuria following eradication, n equals 268) were randomized (1:1) to receive either 875 mg/125mg tablets of AUGMENTIN every 12 hours (n equals 308) or 500 mg/125 mg tablets of AUGMENTIN every 8 hours (n equals 321).

The number of bacteriologically evaluable patients was comparable between the two dosing regimens. AUGMENTIN produced comparable bacteriological success rates in patients assessed 2 to 4 days immediately following end of therapy. The bacteriologic efficacy rates were comparable at one of the follow-up visits (5 to 9 days post-therapy) and at a late post-therapy visit (in the majority of cases, this was 2 to 4 weeks post-therapy), as seen in Table 8.

Table 8: Bacteriologic efficacy rates for AUGMENTIN

Time Post Therapy	875 mg every 12 hours	500 mg every 8 hours
	% (n)	% (n)
2 to 4 days	81% (58)	80% (54)
5 to 9 days	58% (41)	52% (52)
2 to 4 weeks	52% (101)	55% (104)

As noted before, though there was no significant difference in the percentage of adverse events in each group, there was a statistically significant difference in rates of severe diarrhea or withdrawals with diarrhea between the regimens.

14.2 Acute Bacterial Otitis Media and Diarrhea in Pediatric Patients

One US/Canadian clinical trial was conducted which compared 45/6.4 mg/kg/day (divided every 12 hours) of AUGMENTIN for 10 days versus 40/10 mg/kg/day (divided every 8 hours) of AUGMENTIN for 10 days in the treatment of acute otitis media. Only the suspension formulations were used in this trial. A total of 575 pediatric patients (aged 2 months to 12 years) were enrolled, with an even distribution among the 2 treatment groups and a comparable number of patients were evaluable (i.e., greater than or equal to 84%) per treatment group. Otitis media-specific criteria were required for eligibility and a strong correlation was found at the end of therapy and follow-up between these criteria and physician assessment of clinical response. The clinical efficacy rates at the end of therapy visit (defined as 2 to 4 days after the completion of therapy) and at the follow-up visit (defined as 22 to 28 days post-completion of therapy) were comparable for the 2 treatment groups, with the following cure rates obtained for the evaluable patients: At end of therapy, 87% (n equals 265) and 82% (n equals 260) for 45 mg/kg/day every 12 hours and 40 mg/kg/day every 8 hours, respectively. At follow-up, 67% (n equals 249) and 69% (n equals 243) for 45 mg/kg/day every 12 hours and 40 mg/kg/day every 8 hours, respectively.

Diarrhea was defined as either: (a) 3 or more watery or 4 or more loose/watery stools in 1 day; OR (b) 2 watery stools per day or 3 loose/watery stools per day for 2 consecutive days. The incidence of diarrhea was significantly lower in patients who received the every 12 hours regimen compared to patients who received the every 8 hours regimen (14% and 34%, respectively). In addition, the number of patients with either severe diarrhea or who were withdrawn with diarrhea was significantly lower in the every 12 hours treatment group (3% and 8% for the every 12 hours/10 day and every 8 hours/10 day, respectively). In the every 12 hours treatment group, 3 patients (1%) were withdrawn with an allergic reaction, while 1 patient in the every 8 hours group was withdrawn for this reason. The number of patients with a candidal infection of the diaper area was 4% and 6% for the every 12 hours and every 8 hours groups, respectively.

It is not known if the finding of a statistically significant reduction in diarrhea with the oral suspensions dosed every 12 hours, versus suspensions dosed every 8 hours of AUGMENTIN, can be extrapolated to the chewable tablets. The presence of mannitol in the chewable tablets of AUGMENTIN may contribute to a different diarrhea profile. The every 12 hour oral suspensions (200 mg/28.5 mg per 5 mL and 400 mg/57 mg per 5 mL) of AUGMENTIN are sweetened with aspartame.

15 REFERENCES

1. Swanson-Biearman B, Dean BS, Lopez G, Krenzelok EP. The effects of penicillin and cephalosporin ingestions in children less than six years of age. *Vet Hum Toxicol.* 1988; 30: 66-67.

16 HOW SUPPLIED/STORAGE AND HANDLING

AUGMENTIN for Oral Suspension, USP:

125 mg/31.25 mg per 5 mL: Banana-flavored powder for oral suspension, each 5 mL of reconstituted suspension contains 125 mg of amoxicillin as the trihydrate and 31.25 mg of clavulanic acid as the potassium salt (equivalent to 37.23 mg of clavulanate potassium).

NDC71297-240-34	75 mL bottle
NDC 71297-240-35	100 mL bottle
NDC 71297-240-36	150 mL bottle

Dispense in original container.

Store dry powder at or below 25°C (77°F).

Store reconstituted suspension under refrigeration. Discard unused suspension after 10 days.

Keep out of the reach of children.

17 PATIENT COUNSELING INFORMATION

Administration Instructions

Inform patients that AUGMENTIN may be taken every 8 hours or every 12 hours, depending on the dose prescribed. Each dose should be taken with a meal or snack to reduce the possibility of gastrointestinal upset.

Allergic Reactions

Counsel patients that AUGMENTIN contains a penicillin class drug product that can cause allergic reactions in some individuals.

Severe Cutaneous Adverse Reactions (SCAR)

Advise patients about the signs and symptoms of serious skin manifestations. Instruct patients to stop taking AUGMENTIN immediately and promptly report the first signs or symptoms of skin rash, mucosal lesions, or any other sign of hypersensitivity [see *Warnings and Precautions* (5.2)] .

Diarrhea

Counsel patients that diarrhea is a common problem caused by antibacterials, and it usually ends when the antibacterial is discontinued. Sometimes after starting treatment with antibacterials, patients can develop watery and bloody stools (with or without stomach cramps and fever) even as late as 2 or more months after having taken their last dose of the antibacterial. If diarrhea is severe or lasts more than 2 or 3 days, patients should contact their physician as soon as possible.

Antibacterial Resistance

Patients should be counseled that antibacterial drugs, including AUGMENTIN, should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold).

When AUGMENTIN is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may: (1) decrease the effectiveness of the immediate treatment, and (2) increase the likelihood that bacteria will develop resistance and will not be treatable by AUGMENTIN or other antibacterial drugs in the future.

Storage Instructions

Advise patients to keep suspension refrigerated. Shake well before using. When dosing a child with the suspension (liquid) of AUGMENTIN, use a calibrated oral syringe. Be sure to rinse the calibrated oral syringe after each use. Bottles of suspension of AUGMENTIN may contain more liquid than required. Follow your doctor's instructions about the amount to use and the days of treatment your child requires. Discard any unused medicine.

Manufactured For:

Allegis Holdings, LLC
Canton, MS 39046 (USA)

Made in USA

AUGMENTIN is a registered trademark of GlaxoSmithKline and is sub-licensed to Allegis Holdings, LLC.

CLINITEST is a registered trademark of Miles, Inc.

Bottle Label (100 mL) 125mg/31.25mg for Oral Suspension

NDC 71297- 240-35

125 mg/31.25 mg per 5 mL

AUGMENTIN *for oral suspension, USP*

When reconstituted, each 5 mL contains:

AMOXICILLIN, 125 mg as the trihydrate, and

CLAVULANIC ACID, 31.25 mg (equivalent to 37.23 mg of clavulanate potassium)

100 mL (when reconstituted)

Rx only



Directions for mixing:

1. Tap bottle until all powder flows freely.
2. Measure **90 mL** of water (total).
3. Add approximately 2/3 of the water to powder. Replace cap; shake **vigorously**.
4. Add rest of water. Replace cap; shake **vigorously**.

Dosage:
See accompanying prescribing information.

Keep tightly closed. Shake well before each use. Must refrigerate suspension. Date reconstituted

____/____/____.
Discard after 10 days. Some color change is normal during dosing period.

NDC 71297-240-35

125 mg/31.25 mg per 5 mL

AUGMENTIN[®]
amoxicillin and clavulanate potassium for oral suspension, USP

When reconstituted, each 5 mL contains:
AMOXICILLIN, 125 mg as the trihydrate, and
CLAVULANIC ACID, 31.25 mg (equivalent to 37.23 mg of clavulanate potassium)

100 mL
(when reconstituted)



Rx only

Use only if inner seal is intact.
Net contents: Equivalent to 2.5 g amoxicillin and 0.625 g clavulanic acid.
Store dry powder at or below 25° C (77° F).
Mfd. for: Allegis Holdings, LLC, Canton, MS 39046
MADE in USA
Rev. 07/2023

720

Bottle Label (150 mL) 125mg/31.25mg for Oral Suspension

NDC **71297-240-36**

125 mg/31.25 mg per 5 mL

AUGMENTIN for oral suspension, USP

When reconstituted, each 5 mL contains:

AMOXICILLIN, 125 mg as the trihydrate, and

CLAVULANIC ACID, 31.25 mg (equivalent to 37.23 mg of clavulanate potassium)

150 mL (when reconstituted)

Rx only



Directions for mixing:

1. Tap bottle until all powder flows freely.
2. Measure **134 mL** of water (total).
3. Add approximately 2/3 of the water to powder. Replace cap; shake **vigorously**.
4. Add rest of water. Replace cap; shake **vigorously**.

Dosage:
See accompanying prescribing information.

Keep tightly closed. Shake well before each use. Must refrigerate suspension. Date reconstituted

____/____/____.
Discard after 10 days. Some color change is normal during dosing period.

NDC 71297-240-36

125 mg/31.25 mg per 5 mL

AUGMENTIN®
amoxicillin and clavulanate potassium for oral suspension, USP

When reconstituted, each 5 mL contains:
AMOXICILLIN, 125 mg as the trihydrate, and
CLAVULANIC ACID, 31.25 mg (equivalent to 37.23 mg of clavulanate potassium)

150 mL
(when reconstituted)



Rx only

Use only if inner seal is intact.
Net contents: Equivalent to 3.75 g amoxicillin and 0.938 g clavulanic acid.
Store dry powder at or below 25° C (77° F).
Mfd. for: Allegis Holdings, LLC, Canton, MS 39046
MADE in USA
Rev. 07/2023

721

Bottle Label (75 mL) 125mg/31.25mg for Oral Suspension

NDC 71297- 240-34

125 mg/31.25 mg per 5 mL

AUGMENTIN for oral suspension, USP

When reconstituted, each 5 mL contains:

AMOXICILLIN, 125 mg as the trihydrate, and

CLAVULANIC ACID, 31.25 mg (equivalent to 37.23 mg of clavulanate potassium)

75 mL(when reconstituted)

Rx only



Directions for mixing:

1. Tap bottle until all powder flows freely.
2. Measure **67 mL** of water (total).
3. Add approximately 2/3 of the water to powder.
Replace cap; shake **vigorously**.
4. Add rest of water. Replace cap; shake **vigorously**.

Dosage: See accompanying prescribing information.

Keep tightly closed. Shake well before each use. Must refrigerate suspension.

Date reconstituted ____/____/____.
Discard after 10 days.
Some color change is normal during dosing period.

NDC 71297-240-34

125 mg/31.25 mg per 5 mL

AUGMENTIN®
amoxicillin and clavulanate potassium for oral suspension, USP

When reconstituted, each 5 mL contains:
AMOXICILLIN, 125 mg
as the trihydrate, and
CLAVULANIC ACID, 31.25 mg
(equivalent to 37.23 mg of clavulanate potassium)

75 mL
(when reconstituted)



Rx only

Use only if inner seal is intact.
Net contents: Equivalent to 1.875 g amoxicillin and 0.469 g clavulanic acid.
Store dry powder at or below 25° C (77° F).
Mfd. for: Allegis Holdings, LLC, Canton, MS 37620
MADE in USA Rev. 05/2024 730
GTIN: 00371297240340

AUGMENTIN

amoxicillin and clavulanate potassium for suspension

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:71297-240
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMOXICILLIN (UNII: 804826J2HU) (AMOXICILLIN ANHYDROUS - UNII:9EM05410Q9)	AMOXICILLIN ANHYDROUS	125 mg in 5 mL
CLAVULANATE POTASSIUM (UNII: Q42OMW3AT8) (CLAVULANIC ACID - UNII:23521W1S24)	CLAVULANIC ACID	31.25 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
MANNITOL (UNII: 3OWL53L36A)	
SUCCINIC ACID (UNII: AB6MNQ6J6L)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	Score
Shape	Size
Flavor	Imprint Code
	BANANA

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71297-240-35	100 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2023	
2	NDC:71297-240-36	150 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2023	
3	NDC:71297-240-34	75 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/20/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA050575	09/01/2023	

Labeler - Allegis Holdings, LLC (080556861)

Revised: 5/2024

Allegis Holdings, LLC