

STOMACH LIL DRUG STORE PRODUCTS- bismuth subsalicylate liquid
STOMACH CIRCLE K- bismuth subsalicylate liquid
Lil' Drug Store Products, Inc

Lil' Drug Store ® Stomach

Drug Facts

Active ingredient (in each 30 mL dose)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- diarrhea
- travelers' diarrhea
- upset stomach due to overindulgence in food and drink, including
 - heartburn
 - gas
 - indigestion
 - nausea
 - fullness
 - belching

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Contains salicylate. Do not take if you are:

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem

- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over:
 - 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/travelers' diarrhea
 - 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
 - do not exceed 8 doses (240 mL) in 24 hours
 - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** potassium 8 mg, sodium 6 mg
- **each 30 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions?

Call toll free **1-877-507-6516** (M-F 8AM-4:30PM CST)

PRINCIPAL DISPLAY PANEL - 237 mL Bottle Label

QUALITY
GUARANTEED

Compare to the Active
Ingredient in Pepto-Bismol®*

Stomach

Bismuth Subsalicylate 525 mg
Upset Stomach Reliever/Antidiarrheal

Relieves

- Nausea • Heartburn • Indigestion
- Upset Stomach • Diarrhea

8 FL OZ
(237 mL)

Lil'
Drug Store®

<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)</p> <p>Directions • mL = milliliter • shake well before using</p> <ul style="list-style-type: none">• measure only with dosing cup provided. Do not use any other dosing device.• keep dosing cup with product• adults and children 12 years and over:<ul style="list-style-type: none">• 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/travelers' diarrhea• 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)• do not exceed 8 doses (240 mL) in 24 hours• use until diarrhea stops but not more than 2 days• children under 12 years of age: ask a doctor• drink plenty of clear fluids to help prevent dehydration caused by diarrhea <p>Other information • each 30 mL contains: potassium 8 mg, sodium 6 mg</p> <ul style="list-style-type: none">• each 30 mL contains: salicylate 206 mg• low sodium • keep tightly closed• protect from freezing • avoid excessive heat (over 104°F or 40°C) <p>Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum</p> <p>Questions? Call toll free 1-877-507-6516 (M-F 8AM-4:30PM CST)</p>	<p>Compare to the Active Ingredient in Pepto-Bismol®*</p> <p>Stomach</p> <p>Bismuth Subsalicylate 525 mg Upset Stomach Reliever/Antidiarrheal</p> <p>Relieves</p> <ul style="list-style-type: none">• Nausea • Heartburn • Indigestion• Upset Stomach • Diarrhea <p>8 FL OZ (237 mL)</p> <p>Lil' DrugStore</p>	<p>TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.</p> <p>Drug Facts</p> <p>Active ingredient (in each 30 mL dose) Bismuth subsalicylate 525 mg...Purpose Upset stomach reliever/Antidiarrheal</p> <p>Uses relieves • diarrhea • travelers' diarrhea</p> <ul style="list-style-type: none">• upset stomach due to overindulgence in food and drink, including• heartburn • gas • indigestion • nausea • fullness • belching <p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Contains salicylate. Do not take if you are:</p> <ul style="list-style-type: none">• allergic to salicylates (including aspirin)• taking other salicylate products <p>Do not use if you have</p> <ul style="list-style-type: none">• an ulcer • a bleeding problem • bloody or black stool <p>Ask a doctor before use if you have • fever • mucus in the stool</p> <p>Ask a doctor or pharmacist before use if you are taking any drug for</p> <ul style="list-style-type: none">• diabetes • gout • arthritis• anticoagulation (thinning the blood) <p>When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.</p> <p>Stop use and ask a doctor if • symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days</p>
---	---	--

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

Product manufactured for:
Lil' Drug Store Products, Inc.
9300 Earhart Lane SW
Cedar Rapids, IA 52404
68554L-US-12-19
PLD-A492A LB007266



Circle K™ Soothe Upset Stomach -PDP/Package

RELIEF OF NAUSEA, HEARTBURN, INDIGESTION, UPSET STOMACH, DIARRHEA

Compare to the active ingredient
in **Pepto-Bismol**®*

[Circle K logo]

soothe

upset stomach

Bismuth Subsalicylate 525 mg

Upset Stomach Reliever/Antidiarrheal

8 fl oz (237 mL)

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions • mL = milliliter • shake well before using

- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over:
 - 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/travelers' diarrhea
 - 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information • each 30 mL contains: potassium 8 mg, sodium 9 mg

- each 30 mL contains: salicylate 206 mg
- low sodium
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions? Call toll free 1-877-507-6516 (M-F 8AM-4:30PM CST)


*This product is not manufactured or distributed by The Procter & Gamble Company, distributor of Pepto-Bismol®.

Product manufactured for: Lil' Drug Store Products, Inc. 9300 Earhart Lane SW Cedar Rapids, IA 52404

Proudly distributed by Circle K Stores Inc. 5554L-US-06-20-3 PLD-C492B LB007945

RELIEF OF NAUSEA, HEARTBURN,
INDIGESTION, UPSET STOMACH, DIARRHEA

Compare to the active ingredient
in **Pepto-Bismol**®*



CIRCLE K™

soothe

upset stomach

Bismuth Subsalicylate 525 mg
Upset Stomach Reliever/Antidiarrheal

8 fl oz (237 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Drug Facts

Active ingredient (in each 30 mL dose) Bismuth subsalicylate 525 mg...
Purpose Upset stomach reliever/Antidiarrheal

Uses relieves • diarrhea • travelers' diarrhea • upset stomach due to overindulgence in food and drink, including • heartburn • gas • indigestion • nausea • fullness • belching

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are:
 • allergic to salicylates (including aspirin)
 • taking other salicylate products

Do not use if you have
 • an ulcer • a bleeding problem • bloody or black stool

Ask a doctor before use if you have • fever • mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for
 • diabetes • gout • arthritis
 • anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if
 • symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs
 • diarrhea lasts more than 2 days

STOMACH LIL DRUG STORE PRODUCTS

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6855
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII: ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6855-4	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	06/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	12/01/2014	

STOMACH CIRCLE K

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5555
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

XANTHAN GUM (UNII: TTV12P4NEE)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5555-4	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/05/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	12/01/2014	

Labeler - Lil' Drug Store Products, Inc (093103646)

Revised: 5/2024

Lil' Drug Store Products, Inc