

**ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Anti-Diarrheal Drug Facts**

**Active ingredient (in each caplet)**

Loperamide HCl 2 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

**Inactive ingredients**

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

**Questions or comments?**

**1-800-719-9260**

**Principal Display Panel**

meijer®

COMPARE TO IMODIUM® A-D ACTIVE INGREDIENT

Anti-Diarrheal

LOPERAMIDE HYDROCHLORIDE TABLETS,


2 mg | ANTI-DIARRHEAL

24 CAPLETS\*

\*CAPSULE-SHAPED TABLETS

Controls the Symptoms of Diarrhea

Actual Size



22462 6E C4

CODE AREA

OPEN OTHER END

**Drug Facts**

**Active ingredient (in each caplet)** Purpose  
Loperamide HCl 2 mg.....Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl  
**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**

- fever ■ mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are taking a prescription drug.**

Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date



**DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN**  
DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

CONVENIENT RECLOSING TAB



**Drug Facts (continued)**

**Inactive ingredients** anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

**Questions or comments?**  
1-800-719-9260

PID 4586160



\*\*This product is not manufactured or distributed by Johnson & Johnson, owner of the registered trademark Imodium®.

**ANTI DIARRHEAL**

loperamide hydrochloride tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-224
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77T135393C) (LOPERAMIDE - UNII: 6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	

### Product Characteristics

<b>Color</b>	GREEN	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	L2
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-224-53	12 in 1 CARTON	03/24/2003	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41250-224-67	1 in 1 CARTON	02/08/2004	08/31/2021
2		48 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-224-82	1 in 1 CARTON	04/01/2015	09/30/2021
3		200 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:41250-224-91	6 in 1 CARTON	03/16/2016	
4		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:41250-224-80	1 in 1 CARTON	04/30/2003	12/19/2016
5		96 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:41250-224-01	2 in 1 CARTON	04/30/2003	12/19/2016
6		96 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:41250-224-62	24 in 1 CARTON	06/19/2024	
7		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

ANDA	ANDA075232	03/05/2003	
------	------------	------------	--

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 6/2024

Meijer Distribution Inc