

STOMACH RELIEF ULTRA STRENGTH- bismuth subsalicylate liquid

Rite Aid Corporation

Drug Facts

Active ingredient (in each 15 mL)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and over:
 - 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
 - 15 mL (1 dose) every 1/2 as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

Other information

- **each 15 mL contains:** sodium 6 mg
- **each 15 mL contains:** salicylate 227 mg
- low sodium
- keep tightly closed

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in **Pepto-Bismol® Ultra***

2X STRENGTH PER OUNCE†

ULTRA STOMACH RELIEF

BISMUTH SUBSALICYLATE 525 mg

UPSET STOMACH RELIEVER

ANTIDIARRHEAL

Relieves:

heartburn • indigestion • nausea

upset stomach • diarrhea

alcohol free • sugar free

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE

CAMP HILL, PA 17011

www.riteaid.com

Package Label

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ do not take more than 8 doses (120 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ mL = milliliter ■ shake well before using ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 15 mL (1 dose) every ½ hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 15 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ children under 12 years of age: ask a doctor

Other information ■ each 15 mL contains: sodium 5 mg ■ each 15 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company. vs. Original Pepto-Bismol®

DISTRIBUTED BY: RITE AID, 30 HUNTER LANE
CAMP HILL, PA 17011 www.riteaid.com

SATISFACTION GUARANTEE:
If you're not satisfied, we'll happily refund your money.

PLD-B494A
LB007403



2X STRENGTH PER OUNCE!

Compare to the active ingredient in Pepto-Bismol® Ultra*

NDC 11822-7821-5

ULTRA STOMACH RELIEF

BISMUTH SUBSALICYLATE 525 mg

UPSET STOMACH RELIEVER ANTIDIARRHEAL

relieves:
heartburn • indigestion • nausea
upset stomach • diarrhea

alcohol free • sugar free



8 FL OZ (237 mL)

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Active ingredient (in each 15 mL dose)
Bismuth subsalicylate 525 mg.....Upset stomach reliever/Antidiarrheal

Purpose

Uses relieves
■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ indigestion ■ nausea ■ gas ■ fullness ■ belching

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Contains salicylate. Do not take if you are
■ allergic to salicylates (including aspirin)
■ taking other salicylate products

Do not use if you have
■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have
■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are
taking any drug for ■ diabetes ■ gout ■ arthritis
■ anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if
■ symptoms get worse or last more than 2 days
■ ringing in the ears or loss of hearing occurs
■ diarrhea lasts more than 2 days

RITE AID Ultra Stomach Relief

STOMACH RELIEF ULTRA STRENGTH			
bismuth subsalicylate liquid			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-7821
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	525 mg in 15 mL	
Inactive Ingredients			
Ingredient Name	Strength		
BENZOIC ACID (UNII: 8SKN0B0MIM)			
D&C RED NO. 22 (UNII: 1678RKX8RT)			
D&C RED NO. 28 (UNII: 767IP0Y5NH)			
WATER (UNII: 059QF0KO0R)			
GLYCERIN (UNII: PDC6A3C0OX)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
XANTHAN GUM (UNII: TTV12P4NEE)			
Packaging			

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-7821-5	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/27/2019	
2	NDC:11822-7821-6	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/27/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	12/27/2019	

Labeler - Rite Aid Corporation (014578892)

Revised: 10/2023

Rite Aid Corporation