

# **DONEPEZIL HYDROCHLORIDE - donepezil hydrochloride tablet, film coated**

## **Cadila Pharmaceuticals Limited**

-----

### **HIGHLIGHTS OF PRESCRIBING INFORMATION**

**These highlights do not include all the information needed to use DONEPEZIL HYDROCHLORIDE TABLETS safely and effectively. See full prescribing information for DONEPEZIL HYDROCHLORIDE TABLETS.**

**DONEPEZIL HYDROCHLORIDE tablets USP, for oral use**

**Initial U.S. Approval: 1996**

### **INDICATIONS AND USAGE**

-----

Donepezil Hydrochloride is an acetylcholinesterase inhibitor indicated for the treatment of dementia of the Alzheimer's type. Efficacy has been demonstrated in patients with mild, moderate, and severe Alzheimer's Disease.(1)

### **DOSAGE AND ADMINISTRATION**

-----

- Mild to Moderate Alzheimer's Disease: 5 mg to 10 mg once daily (2.1)
- Moderate to Severe Alzheimer's Disease: 10 mg once daily (2.2)

### **DOSAGE FORMS AND STRENGTHS**

-----

- Tablets: 5 mg and 10 mg (3)

### **CONTRAINDICATIONS**

-----

- Known hypersensitivity to donepezil hydrochloride or to piperidine derivatives (4)

### **WARNINGS AND PRECAUTIONS**

-----

- Cholinesterase inhibitors are likely to exaggerate succinylcholine-type muscle relaxation during anesthesia (5.1)
- Cholinesterase inhibitors may have vagotonic effects on the sinoatrial and atrioventricular nodes manifesting as bradycardia or heart block (5.2)
- Donepezil Hydrochloride can cause vomiting. Patients should be observed closely at initiation of treatment and after dose increases (5.3)
- Patients should be monitored closely for symptoms of active or occult gastrointestinal (GI) bleeding, especially those at increased risk for developing ulcers (5.4)
- Cholinomimetics may cause bladder outflow obstructions (5.6)
- Cholinomimetics are believed to have some potential to cause generalized convulsions (5.7)
- Cholinesterase inhibitors should be prescribed with care to patients with a history of asthma or obstructive pulmonary disease (5.8)

### **ADVERSE REACTIONS**

-----

Most common adverse reactions in clinical studies of Donepezil Hydrochloride Tablets are nausea, diarrhea, insomnia, vomiting, muscle cramps, fatigue, and anorexia (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Cadila Pharmaceuticals Limited at 1-202-355-9785 (fax 1-202-355-9784) or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### **DRUG INTERACTIONS**

-----

- Cholinesterase inhibitors have the potential to interfere with the activity of anticholinergic medications (7.1)
- A synergistic effect may be expected with concomitant administration of succinylcholine, similar neuromuscular blocking agents, or cholinergic agonists (7.2)

### **USE IN SPECIFIC POPULATIONS**

-----

Pregnancy: Based on animal data, may cause fetal harm (8.1)

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Dosing in Mild to Moderate Alzheimer's Disease
- 2.2 Dosing in Moderate to Severe Alzheimer's Disease
- 2.3 Administration Information

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Anesthesia
- 5.2 Cardiovascular Conditions
- 5.3 Nausea and Vomiting
- 5.4 Peptic Ulcer Disease and GI Bleeding
- 5.5 Weight Loss
- 5.6 Genitourinary Conditions
- 5.7 Neurological Conditions: Seizures
- 5.8 Pulmonary Conditions

**6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

- 7.1 Use with Anticholinergics
- 7.2 Use with Cholinomimetics and Other Cholinesterase Inhibitors

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Lower Weight Individuals

**10 OVERDOSAGE**

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

**13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
- 13.2 Animal Toxicology and/or Pharmacology

**14 CLINICAL STUDIES**

- 14.1 Mild to Moderate Alzheimer's Disease
- 14.2 Moderate to Severe Alzheimer's Disease

**16 HOW SUPPLIED/STORAGE AND HANDLING**

- 16.1 Donepezil Hydrochloride Tablets

**17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

Donepezil Hydrochloride Tablets is indicated for the treatment of dementia of the Alzheimer's type. Efficacy has been demonstrated in patients with mild, moderate, and severe Alzheimer's disease.

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Dosing in Mild to Moderate Alzheimer's Disease**

The recommended starting dosage of Donepezil Hydrochloride Tablets is 5 mg administered once per day in the evening, just prior to retiring. The maximum recommended dosage of Donepezil Hydrochloride Tablets in patients with mild to moderate Alzheimer's disease is 10 mg per day. A dose of 10 mg should not be administered until patients have been on a daily dose of 5 mg for 4 to 6 weeks.

#### **2.2 Dosing in Moderate to Severe Alzheimer's Disease**

The recommended starting dosage of Donepezil Hydrochloride Tablets is 5 mg administered once per day in the evening, just prior to retiring. The maximum recommended dosage of Donepezil Hydrochloride Tablets in patients with moderate to severe Alzheimer's disease is 23 mg per day. A dose of 10 mg should not be administered until patients have been on a daily dose of 5 mg for 4 to 6 weeks.

#### **2.3 Administration Information**

Donepezil Hydrochloride Tablets should be taken in the evening, just prior to retiring. Donepezil Hydrochloride Tablets can be taken with or without food.

### **3 DOSAGE FORMS AND STRENGTHS**

Donepezil Hydrochloride Tablets is supplied as film-coated, round tablets containing 5 mg and 10 mg of donepezil hydrochloride.

- The 5 mg tablets are white, round, biconvex film coated tablets, debossed 'C57' on one side and plain on other side.
- The 10 mg tablets are yellow to light yellow, round, biconvex film coated tablets, debossed 'C55' on one side and plain on other side.

## **4 CONTRAINDICATIONS**

Donepezil Hydrochloride Tablets is contraindicated in patients with known hypersensitivity to donepezil hydrochloride or to piperidine derivatives.

## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Anesthesia**

Donepezil Hydrochloride Tablets, as a cholinesterase inhibitor, is likely to exaggerate succinylcholine-type muscle relaxation during anesthesia.

### **5.2 Cardiovascular Conditions**

Because of their pharmacological action, cholinesterase inhibitors may have vagotonic effects on the sinoatrial and atrioventricular nodes. This effect may manifest as bradycardia or heart block in patients both with and without known underlying cardiac conduction abnormalities. Syncopal episodes have been reported in association with the use of Donepezil Hydrochloride Tablets.

### **5.3 Nausea and Vomiting**

Donepezil Hydrochloride Tablets, as a predictable consequence of its pharmacological properties, has been shown to produce diarrhea, nausea, and vomiting. These effects, when they occur, appear more frequently with the 10 mg/day dose than with the 5 mg/day dose.

Although in most cases, these effects have been transient, sometimes lasting one to three weeks, and have resolved during continued use of Donepezil Hydrochloride Tablets, patients should be observed closely at the initiation of treatment and after dose increases.

### **5.4 Peptic Ulcer Disease and GI Bleeding**

Through their primary action, cholinesterase inhibitors may be expected to increase gastric acid secretion due to increased cholinergic activity. Therefore, patients should be monitored closely for symptoms of active or occult gastrointestinal bleeding, especially those at increased risk for developing ulcers, e.g., those with a history of ulcer disease or those receiving concurrent nonsteroidal anti-inflammatory drugs (NSAIDs). Clinical studies of Donepezil Hydrochloride Tablets in a dose of 5 mg/day to 10 mg/day have shown no increase, relative to placebo, in the incidence of either peptic ulcer disease or gastrointestinal bleeding. Results of a controlled clinical study with 23 mg/day showed an increase, relative to 10 mg/day, in the incidence of peptic ulcer disease (0.4% vs. 0.2%) and gastrointestinal bleeding from any site (1.1% vs. 0.6%).

### **5.5 Weight Loss**

Weight loss was reported as an adverse event in 4.7% of patients assigned to Donepezil Hydrochloride Tablets in a dose of 23 mg/day compared to 2.5% of patients assigned to 10 mg/day. Compared to their baseline weights, 8.4% of patients taking 23 mg/day were found to have a weight decrease of  $\geq 7\%$  by the end of the study, while 4.9% of patients taking 10 mg/day were found to have weight loss of  $\geq 7\%$  at the end of the study.

## 5.6 Genitourinary Conditions

Although not observed in clinical trials of Donepezil Hydrochloride Tablets, cholinomimetics may cause bladder outflow obstruction.

## 5.7 Neurological Conditions: Seizures

Cholinomimetics are believed to have some potential to cause generalized convulsions. However, seizure activity also may be a manifestation of Alzheimer's disease.

## 5.8 Pulmonary Conditions

Because of their cholinomimetic actions, cholinesterase inhibitors should be prescribed with care to patients with a history of asthma or obstructive pulmonary disease.

# 6 ADVERSE REACTIONS

The following serious adverse reactions are described below and elsewhere in the labeling:

- Cardiovascular Conditions *[see Warnings and Precautions (5.2)]*
- Nausea and Vomiting *[see Warnings and Precautions (5.3)]*
- Peptic Ulcer Disease and GI Bleeding *[see Warnings and Precautions (5.4)]*
- Weight Loss *[see Warnings and Precautions (5.5)]*
- Genitourinary Conditions *[see Warnings and Precautions (5.6)]*
- Neurological Conditions: Seizures *[see Warnings and Precautions (5.7)]*
- Pulmonary Conditions *[see Warnings and Precautions (5.8)]*

## 6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Donepezil Hydrochloride Tablets has been administered to over 1,700 individuals during clinical trials worldwide. Approximately 1,200 of these patients have been treated for at least 3 months and more than 1,000 patients have been treated for at least 6 months.

Controlled and uncontrolled trials in the United States included approximately 900 patients. In regards to the highest dose of 10 mg/day, this population includes 650 patients treated for 3 months, 475 patients treated for 6 months, and 116 patients treated for over 1 year. The range of patient exposure is from 1 to 1,214 days.

### Mild to Moderate Alzheimer’s Disease

#### *Adverse Reactions Leading to Discontinuation*

The rates of discontinuation from controlled clinical trials of Donepezil Hydrochloride Tablets due to adverse reactions for the Donepezil Hydrochloride Tablets 5 mg/day treatment groups were comparable to those of placebo treatment groups at approximately 5%. The rate of discontinuation of patients who received 7-day escalation from 5 mg/day to 10 mg/day was higher at 13%.

The most common adverse reactions leading to discontinuation, defined as those occurring in at least 2% of patients and at twice or more the incidence seen in placebo patients, are shown in Table 1.

**Table 1. Most Common Adverse Reactions Leading to Discontinuation in Patients with Mild to Moderate Alzheimer’s Disease**

<b>Adverse Reaction</b>	<b>Placebo (n=355) %</b>	<b>5 mg/day Donepezil Hydrochloride Tablets (n=350) %</b>	<b>10 mg/day Donepezil Hydrochloride Tablets (n=315) %</b>
Nausea	1	1	3
Diarrhea	0	<1	3
Vomiting	<1	<1	2

#### *Most Common Adverse Reactions*

The most common adverse reactions, defined as those occurring at a frequency of at least 5% in patients receiving 10 mg/day and twice the placebo rate, are largely predicted by Donepezil Hydrochloride Tablet's cholinomimetic effects. These include nausea, diarrhea, insomnia, vomiting, muscle cramp, fatigue and anorexia. These adverse reactions were often transient, resolving during continued Donepezil Hydrochloride Tablets treatment without the need for dose modification.

There is evidence to suggest that the frequency of these common adverse reactions

may be affected by the rate of titration. An open-label study was conducted with 269 patients who received placebo in the 15- and 30-week studies. These patients were titrated to a dose of 10 mg/day over a 6-week period. The rates of common adverse reactions were lower than those seen in patients titrated to 10 mg/day over one week in the controlled clinical trials and were comparable to those seen in patients on 5 mg/day.

See Table 2 for a comparison of the most common adverse reactions following one and six week titration regimens.

**Table 2. Comparison of Rates of Adverse Reactions in Mild to Moderate Patients Titrated to 10 mg/day over 1 and 6 Weeks**

Adverse Reaction	No titration		One week titration	Six week titration
	Placebo (n=315) %	5 mg/day (n=311) %	10 mg/day (n=315) %	10 mg/day (n=269) %
Nausea	6	5	19	6
Diarrhea	5	8	15	9
Insomnia	6	6	14	6
Fatigue	3	4	8	3
Vomiting	3	3	8	5
Muscle cramps	2	6	8	3
Anorexia	2	3	7	3

Table 3 lists adverse reactions that occurred in at least 2% of patients in pooled placebo-controlled trials who received either Donepezil Hydrochloride Tablets 5 mg or 10 mg and for which the rate of occurrence was greater for patients treated with Donepezil Hydrochloride Tablets than with placebo. In general, adverse reactions occurred more frequently in female patients and with advancing age.

**Table 3. Adverse Reactions in Pooled Placebo-Controlled Clinical Trials in Mild to Moderate Alzheimer's Disease**

Adverse Reaction	Placebo (n=355) %	Donepezil Hydrochloride Tablets (n=747) %
<b>Percent of Patients with any Adverse Reaction</b>	<b>72</b>	<b>74</b>
Nausea	6	11
Diarrhea	5	10

Headache	9	10
Insomnia	6	9
Pain, various locations	8	9
Dizziness	6	8
Accident	6	7
Muscle Cramps	2	6
Fatigue	3	5
Vomiting	3	5
Anorexia	2	4
Ecchymosis	3	4
Abnormal Dreams	0	3
Depression	< 1	3
Weight Loss	1	3
Arthritis	1	2
Frequent Urination	1	2
Somnolence	< 1	2
Syncope	1	2

### Severe Alzheimer's Disease (Donepezil Hydrochloride Tablets 5 mg/day and 10 mg/day)

Donepezil Hydrochloride Tablets has been administered to over 600 patients with severe Alzheimer's disease during clinical trials of at least 6 months duration, including three double-blind, placebo-controlled trials, two of which had an open label extension.

#### *Adverse Reactions Leading to Discontinuation*

The rates of discontinuation from controlled clinical trials of Donepezil Hydrochloride Tablets due to adverse reactions for the Donepezil Hydrochloride Tablets patients were approximately 12% compared to 7% for placebo patients. The most common adverse reactions leading to discontinuation, defined as those occurring in at least 2% of Donepezil Hydrochloride Tablets patients and at twice or more the incidence seen in placebo, were anorexia (2% vs. 1% placebo), nausea (2% vs. <1% placebo), diarrhea (2% vs. 0% placebo), and urinary tract infection (2% vs. 1% placebo).

#### *Most Common Adverse Reactions*

The most common adverse reactions, defined as those occurring at a frequency of at least 5% in patients receiving Donepezil Hydrochloride Tablets and at twice or more the placebo rate, are largely predicted by Donepezil Hydrochloride Tablet's cholinomimetic effects. These include diarrhea, anorexia, vomiting, nausea, and ecchymosis. These adverse reactions were often transient, resolving during continued Donepezil Hydrochloride Tablets treatment without the need for dose modification.

Table 4 lists adverse reactions that occurred in at least 2% of patients in pooled placebo-controlled trials who received Donepezil Hydrochloride Tablets 5 mg or 10 mg and for which the rate of occurrence was greater for patients treated with Donepezil Hydrochloride Tablets than with placebo.



**Table 4. Adverse Reactions in Pooled Controlled Clinical Trials in Severe Alzheimer's Disease**

<b>Body System/Adverse Reaction</b>	<b>Placebo (n=392) %</b>	<b>Donepezil Hydrochloride Tablets (n=501) %</b>
<b>Percent of Patients with any Adverse Reaction</b>	<b>73</b>	<b>81</b>
Accident	12	13
Infection	9	11
Diarrhea	4	10
Anorexia	4	8
Vomiting	4	8
Nausea	2	6
Insomnia	4	5
Ecchymosis	2	5
Headache	3	4
Hypertension	2	3
Pain	2	3
Back Pain	2	3
Eczema	2	3
Hallucinations	1	3
Hostility	2	3
Increase in Creatine Phosphokinase	1	3
Nervousness	2	3
Fever	1	2
Chest Pain	<1	2
Confusion	1	2
Dehydration	1	2
Depression	1	2
Dizziness	1	2
Emotional Lability	1	2
Hemorrhage	1	2
Hyperlipemia	<1	2
Personality Disorder	1	2
Somnolence	1	2
Syncope	1	2
Urinary Incontinence	1	2

Moderate to Severe Alzheimer's Disease (Donepezil Hydrochloride Tablets 23 mg/day)  
 Donepezil Hydrochloride Tablets 23 mg/day has been administered to over 1,300 individuals globally in clinical trials. Approximately 1,050 of these patients have been treated for at least three months and more than 950 patients have been treated for at least six months. The range of patient exposure was from 1 to over 500 days.

### *Adverse Reactions Leading to Discontinuation*

The rate of discontinuation from a controlled clinical trial of Donepezil Hydrochloride Tablets 23 mg/day due to adverse reactions was higher (19%) than for the 10 mg/day treatment group (8%). The most common adverse reactions leading to discontinuation, defined as those occurring in at least 1% of patients and greater than those occurring with 10 mg/day are shown in Table 5.

**Table 5. Most Common Adverse Reactions Leading to Discontinuation in Patients with Moderate to Severe Alzheimer's Disease**

<b>Adverse Reaction</b>	<b>23 mg/day Donepezil Hydrochloride Tablets (n=963) %</b>	<b>10 mg/day Donepezil Hydrochloride Tablets (n=471) %</b>
Vomiting	3	0
Diarrhea	2	0
Nausea	2	0
Dizziness	1	0

The majority of discontinuations due to adverse reactions in the 23 mg group occurred during the first month of treatment.

### *Most Common Adverse Reactions with Donepezil Hydrochloride Tablets 23 mg/day.*

The most common adverse reactions, defined as those occurring at a frequency of at least 5%, include nausea, diarrhea, vomiting, and anorexia.

Table 6 lists adverse reactions that occurred in at least 2% of patients who received 23 mg/day of Donepezil Hydrochloride Tablets and at a higher frequency than those receiving 10 mg/day of Donepezil Hydrochloride Tablets in a controlled clinical trial that compared the two doses. In this study, there were no important differences in the type of adverse reactions in patients taking Donepezil Hydrochloride Tablets with or without memantine.

**Table 6. Adverse Reactions in a Controlled Clinical Trial in Moderate to Severe Alzheimer's Disease**

<b>Adverse Reaction</b>	<b>23 mg/day Donepezil Hydrochloride Tablets</b>	<b>10 mg/day Donepezil Hydrochloride Tablets</b>
-------------------------	--	--

	(n=963) %	(n=471) %
<b>Percent of Patients with any Adverse Reaction</b>	74	64
Nausea	12	3
Vomiting	9	3
Diarrhea	8	5
Anorexia	5	2
Dizziness	5	3
Weight Loss	5	3
Headache	4	3
Insomnia	3	2
Urinary incontinence	3	1
Asthenia	2	1
Contusion	2	0
Fatigue	2	1
Somnolence	2	1

## 6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of Donepezil Hydrochloride Tablets. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Abdominal pain, agitation, aggression, cholecystitis, confusion, convulsions, hallucinations, heart block (all types), hemolytic anemia, hepatitis, hyponatremia, neuroleptic malignant syndrome, pancreatitis, rash, rhabdomyolysis, QTc prolongation, and torsade de pointes.

## 7 DRUG INTERACTIONS

### 7.1 Use with Anticholinergics

Because of their mechanism of action, cholinesterase inhibitors have the potential to interfere with the activity of anticholinergic medications.

### 7.2 Use with Cholinomimetics and Other Cholinesterase Inhibitors

A synergistic effect may be expected when cholinesterase inhibitors are given concurrently with succinylcholine, similar neuromuscular blocking agents or cholinergic agonists such as bethanechol.

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

### Risk Summary

There are no adequate data on the developmental risks associated with the use of Donepezil Hydrochloride Tablets in pregnant women. In animal studies, developmental toxicity was not observed when donepezil was administered to pregnant rats and rabbits during organogenesis, but administration to rats during the latter part of pregnancy and throughout lactation resulted in increased stillbirths and decreased offspring survival at clinically relevant doses [see *Data*]. In the U.S. general population, the estimated background risks of major birth defects and miscarriage in clinically recognized pregnancies are 2% to 4% and 15% to 20%, respectively. The background risks of major birth defects and miscarriage for the indicated population are unknown.

### Data

#### *Animal Data*

Oral administration of donepezil to pregnant rats and rabbits during the period of organogenesis did not produce any teratogenic effects at doses up to 16 mg/kg/day (approximately 6 times the maximum recommended human dose [MRHD] of 23 mg/day on a mg/m<sup>2</sup> basis) and 10 mg/kg/day (approximately 7 times the MRHD on a mg/m<sup>2</sup> basis), respectively. Oral administration of donepezil (1, 3, 10 mg/kg/day) to rats during late gestation and throughout lactation to weaning produced an increase in stillbirths and reduced offspring survival through postpartum day 4 at the highest dose. The no-effect dose of 3 mg/kg/day is approximately equal to the MRHD on a mg/m<sup>2</sup> basis.

## **8.2 Lactation**

### Risk Summary

There are no data on the presence of donepezil or its metabolites in human milk, the effects on the breastfed infant, or on milk production

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Donepezil Hydrochloride Tablets and any potential adverse effects on the breastfed infant from Donepezil Hydrochloride Tablets or from the underlying maternal condition.

## **8.4 Pediatric Use**

The safety and effectiveness in pediatric patients have not been established.

## **8.5 Geriatric Use**

Alzheimer's disease is a disorder occurring primarily in individuals over 55 years of age. The mean age of patients enrolled in the clinical studies with Donepezil Hydrochloride Tablets was 73 years; 80% of these patients were between 65 and 84 years old, and 49% of patients were at or above the age of 75. The efficacy and safety data presented in the clinical trials section were obtained from these patients. There were no clinically significant differences in most adverse reactions reported by patient groups  $\geq$  65 years old and  $<$  65 years old.

## 8.6 Lower Weight Individuals

In the controlled clinical trial, among patients in the Donepezil Hydrochloride Tablets 23 mg treatment group, those patients weighing < 55 kg reported more nausea, vomiting, and decreased weight than patients weighing 55 kg or more. There were more withdrawals due to adverse reactions as well. This finding may be related to higher plasma exposure associated with lower weight.

## 10 OVERDOSAGE


Because strategies for the management of overdose are continually evolving, it is advisable to contact a PoisonControlCenter to determine the latest recommendations for the management of an overdose of any drug.

As in any case of overdose, general supportive measures should be utilized. Overdosage with cholinesterase inhibitors can result in cholinergic crisis characterized by severe nausea, vomiting, salivation, sweating, bradycardia, hypotension, respiratory depression, collapse and convulsions. Increasing muscle weakness is a possibility and may result in death if respiratory muscles are involved. Tertiary anticholinergics such as atropine may be used as an antidote for Donepezil Hydrochloride Tablets overdose. Intravenous atropine sulfate titrated to effect is recommended: an initial dose of 1.0 to 2.0 mg IV with subsequent doses based upon clinical response. Atypical responses in blood pressure and heart rate have been reported with other cholinomimetics when co-administered with quaternary anticholinergics such as glycopyrrolate. It is not known whether Donepezil Hydrochloride Tablets and/or its metabolites can be removed by dialysis (hemodialysis, peritoneal dialysis, or hemofiltration).

Dose-related signs of toxicity in animals included reduced spontaneous movement, prone position, staggering gait, lacrimation, clonic convulsions, depressed respiration, salivation, miosis, tremors, fasciculation and lower body surface temperature.

## 11 DESCRIPTION

Donepezil Hydrochloride is a reversible inhibitor of the enzyme acetylcholinesterase, known chemically as (±)-2, 3-dihydro-5, 6-dimethoxy-2-[[1-(phenylmethyl)-4-piperidinyl]methyl]-1*H*-inden-1-one hydrochloride. Donepezil hydrochloride is commonly referred to in the pharmacological literature as E2020. It has an empirical formula of C<sub>24</sub>H<sub>29</sub>NO<sub>3</sub>HCl and a molecular weight of 415.96. Donepezil hydrochloride is a white crystalline powder and is freely soluble in chloroform, soluble in water and in glacial acetic acid, slightly soluble in ethanol and in acetonitrile and practically insoluble in ethyl acetate and in n-hexane.



Donepezil Hydrochloride Tablets is available for oral administration in film-coated tablets containing 5 mg or 10 mg of donepezil hydrochloride.

Inactive ingredients in 5 mg and 10 mg tablets are lactose monohydrate, microcrystalline cellulose, pre-gelatinized starch, hydroxypropyl cellulose and magnesium stearate. The film coating contains hypromellose, polyethylene glycol, talc, titanium dioxide. Additionally, the 10 mg tablet contains yellow iron oxide as a coloring agent. Organic Impurities are analyzed as per procedure 2 of USP monograph.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Current theories on the pathogenesis of the cognitive signs and symptoms of Alzheimer's disease attribute some of them to a deficiency of cholinergic neurotransmission.

Donepezil hydrochloride is postulated to exert its therapeutic effect by enhancing cholinergic function. This is accomplished by increasing the concentration of acetylcholine through reversible inhibition of its hydrolysis by acetylcholinesterase. There is no evidence that donepezil alters the course of the underlying dementing process.

### 12.3 Pharmacokinetics

Pharmacokinetics of donepezil are linear over a dose range of 1-10 mg given once daily. The rate and extent of absorption of Donepezil Hydrochloride Tablets are not influenced by food.

Based on population pharmacokinetic analysis of plasma donepezil concentrations measured in patients with Alzheimer's disease, following oral dosing, peak plasma concentration is achieved for Donepezil Hydrochloride 23 mg tablets in approximately 8 hours, compared with 3 hours for Donepezil Hydrochloride 10 mg tablets. Peak plasma concentrations were about 2-fold higher for Donepezil Hydrochloride 23 mg tablets than Donepezil Hydrochloride 10 mg tablets.

The elimination half life of donepezil is about 70 hours, and the mean apparent plasma clearance (Cl/F) is 0.13 - 0.19 L/hr/kg. Following multiple dose administration, donepezil accumulates in plasma by 4-7 fold, and steady state is reached within 15 days. The steady state volume of distribution is 12 - 16 L/kg. Donepezil is approximately 96%

bound to human plasma proteins, mainly to albumins (about 75%) and alpha<sub>1</sub> - acid glycoprotein (about 21%) over the concentration range of 2-1,000 ng/mL.

Donepezil is both excreted in the urine intact and extensively metabolized to four major metabolites, two of which are known to be active, and a number of minor metabolites, not all of which have been identified. Donepezil is metabolized by CYP 450 isoenzymes 2D6 and 3A4 and undergoes glucuronidation. Following administration of <sup>14</sup>C-labeled donepezil, plasma radioactivity, expressed as a percent of the administered dose, was present primarily as intact donepezil (53%) and as 6-O-desmethyl donepezil (11%), which has been reported to inhibit AChE to the same extent as donepezil *in vitro* and was found in plasma at concentrations equal to about 20% of donepezil. Approximately 57% and 15% of the total radioactivity was recovered in urine and feces, respectively, over a period of 10 days, while 28% remained unrecovered, with about 17% of the donepezil dose recovered in the urine as unchanged drug. Examination of the effect of CYP2D6 genotype in Alzheimer's patients showed differences in clearance values among CYP2D6 genotype subgroups. When compared to the extensive metabolizers, poor metabolizers had a 31.5% slower clearance and ultra-rapid metabolizers had a 24% faster clearance.

#### Hepatic Disease

In a study of 10 patients with stable alcoholic cirrhosis, the clearance of Donepezil hydrochloride Tablets was decreased by 20% relative to 10 healthy age- and sex-matched subjects.

#### Renal Disease

In a study of 11 patients with moderate to severe renal impairment ( $Cl_c < 18$  mL/min/1.73 m<sup>2</sup>) the clearance of Donepezil Hydrochloride Tablets did not differ from 11 age- and sex-matched healthy subjects.

#### Age

No formal pharmacokinetic study was conducted to examine age-related differences in the pharmacokinetics of Donepezil Hydrochloride Tablets. Population pharmacokinetic analysis suggested that the clearance of donepezil in patients decreases with increasing age. When compared with 65-year old, subjects, 90-year old subjects have a 17% decrease in clearance, while 40-year old subjects have a 33% increase in clearance. The effect of age on donepezil clearance may not be clinically significant.

#### Gender and Race

No specific pharmacokinetic study was conducted to investigate the effects of gender and race on the disposition of Donepezil Hydrochloride Tablets. However, retrospective pharmacokinetic analysis and population pharmacokinetic analysis of plasma donepezil concentrations measured in patients with Alzheimer's disease indicates that gender and race (Japanese and Caucasians) did not affect the clearance of Donepezil Hydrochloride Tablets to an important degree.

#### Body Weight

There was a relationship noted between body weight and clearance. Over the range of body weight from 50 kg to 110 kg, clearance increased from 7.77 L/h to 14.04 L/h, with a value of 10 L/hr for 70 kg individuals.

## Drug Interactions

### *Effect of Donepezil Hydrochloride Tablets on the Metabolism of Other Drugs*

No *in vivo* clinical trials have investigated the effect of Donepezil Hydrochloride Tablets on the clearance of drugs metabolized by CYP 3A4 (e.g., cisapride, terfenadine) or by CYP 2D6 (e.g., imipramine). However, *in vitro* studies show a low rate of binding to these enzymes (mean  $K_i$  about 50-130 micrometre), that, given the therapeutic plasma concentrations of donepezil (164 nM), indicates little likelihood of interference. Based on *in vitro* studies, donepezil shows little or no evidence of direct inhibition of CYP2B6, CYP2C8, and CYP2C19 at clinically relevant concentrations.

Whether Donepezil Hydrochloride Tablets has any potential for enzyme induction is not known. Formal pharmacokinetic studies evaluated the potential of Donepezil Hydrochloride Tablets for interaction with theophylline, cimetidine, warfarin, digoxin, and ketoconazole. No effects of Donepezil Hydrochloride Tablets on the pharmacokinetics of these drugs were observed.

### *Effect of Other Drugs on the Metabolism of Donepezil Hydrochloride Tablets*

Ketoconazole and quinidine, strong inhibitors of CYP450 3A and 2D6, respectively, inhibit donepezil metabolism *in vitro*. Whether there is a clinical effect of quinidine is not known. Population pharmacokinetic analysis showed that in the presence of concomitant CYP2D6 inhibitors donepezil AUC was increased by approximately 17% to 20% in Alzheimer's disease patients taking Donepezil Hydrochloride Tablets 10 and 23 mg. This represented an average effect of weak, moderate, and strong CYP2D6 inhibitors. In a 7-day crossover study in 18 healthy volunteers, ketoconazole (200 mg q.d.) increased mean donepezil (5mg q.d.) concentrations ( $AUC_{0-24}$  and  $C_{max}$ ) by 36%. The clinical relevance of this increase in concentration is unknown.

Inducers of CYP 3A (e.g., phenytoin, carbamazepine, dexamethasone, rifampin, and phenobarbital) could increase the rate of elimination of Donepezil Hydrochloride Tablets.

Formal pharmacokinetic studies demonstrated that the metabolism of Donepezil Hydrochloride Tablets is not significantly affected by concurrent administration of digoxin or cimetidine.

An *in vitro* study showed that donepezil was not a substrate of P-glycoprotein.

### *Drugs Highly Bound to Plasma Proteins*

Drug displacement studies have been performed *in vitro* between this highly bound drug (96%) and other drugs such as furosemide, digoxin, and warfarin. Donepezil Hydrochloride Tablets at concentrations of 0.3-10 micrograms/mL did not affect the binding of furosemide (5 micrograms/mL), digoxin (2 ng/mL), and warfarin (3 micrograms/mL) to human albumin. Similarly, the binding of Donepezil Hydrochloride Tablets to human albumin was not affected by furosemide, digoxin and warfarin.

## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**



No evidence of carcinogenic potential was obtained in an 88-week carcinogenicity study of donepezil conducted in mice at oral doses up to 180 mg/kg/day (approximately 40 times the maximum recommended human dose [MRHD] of 23 mg/day on a mg/m<sup>2</sup> basis), or in a 104-week carcinogenicity study in rats at oral doses up to 30 mg/kg/day (approximately 13 times the MRHD on a mg/m<sup>2</sup> basis).

Donepezil was negative in a battery of genotoxicity assays (*in vitro* bacterial reverse mutation, *in vitro* mouse lymphoma *tk*, *in vitro* chromosomal aberration, and *in vivo* mouse micronucleus).

Donepezil had no effect on fertility in rats at oral doses up to 10 mg/kg/day (approximately 4 times the MRHD on a mg/m<sup>2</sup> basis) when administered to males and females prior to and during mating and continuing in females through implantation.

### **13.2 Animal Toxicology and/or Pharmacology**

In an acute dose neurotoxicity study in female rats, oral administration of donepezil and memantine in combination resulted in increased incidence, severity, and distribution of neurodegeneration compared with memantine alone. The no-effect levels of the combination were associated with clinically relevant plasma donepezil and memantine levels.

The relevance of this finding to humans is unknown.

## **14 CLINICAL STUDIES**

### **14.1 Mild to Moderate Alzheimer's Disease**

The effectiveness of Donepezil Hydrochloride Tablets as a treatment for mild to moderate Alzheimer's disease is demonstrated by the results of two randomized, double-blind, placebo-controlled clinical investigations in patients with Alzheimer's disease (diagnosed by NINCDS and DSM III-R criteria, Mini-Mental State Examination  $\geq 10$  and  $\leq 26$  and Clinical Dementia Rating of 1 or 2). The mean age of patients participating in Donepezil Hydrochloride Tablets trials was 73 years with a range of 50 to 94. Approximately 62% of patients were women and 38% were men. The racial distribution was white 95%, black 3% and other races 2%.

The higher dose of 10 mg did not provide a statistically significantly greater clinical benefit than 5 mg. There is a suggestion, however, based upon order of group mean scores and dose trend analyses of data from these clinical trials, that a daily dose of 10 mg of Donepezil Hydrochloride Tablets might provide additional benefit for some patients. Accordingly, whether or not to employ a dose of 10 mg is a matter of prescriber and patient preference.

#### Study Outcome Measures

In each study, the effectiveness of treatment with Donepezil Hydrochloride Tablets was evaluated using a dual outcome assessment strategy.

The ability of Donepezil Hydrochloride Tablets to improve cognitive performance was

assessed with the cognitive subscale of the Alzheimer's Disease Assessment Scale (ADAS-cog), a multi-item instrument that has been extensively validated in longitudinal cohorts of Alzheimer's disease patients. The ADAS-cog examines selected aspects of cognitive performance including elements of memory, orientation, attention, reasoning, language and praxis. The ADAS-cog scoring range is from 0 to 70, with higher scores indicating greater cognitive impairment. Elderly normal adults may score as low as 0 or 1, but it is not unusual for non-demented adults to score slightly higher.

The patients recruited as participants in each study had mean scores on the ADAS-cog of approximately 26 points, with a range from 4 to 61. Experience based on longitudinal studies of ambulatory patients with mild to moderate Alzheimer's disease suggest that scores on the ADAS-cog increase (worsen) by 6 - 12 points per year. However, smaller changes may be seen in patients with very mild or very advanced disease since the ADAS-cog is not uniformly sensitive to change over the course of the disease. The annualized rate of decline in the placebo patients participating in Donepezil Hydrochloride Tablets trials was approximately 2 to 4 points per year.

The ability of Donepezil Hydrochloride Tablets to produce an overall clinical effect was assessed using a Clinician's Interview-Based Impression of Change that required the use of caregiver information, the CIBIC-plus. The CIBIC-plus is not a single instrument and is not a standardized instrument like the ADAS-cog. Clinical trials for investigational drugs have used a variety of CIBIC formats, each different in terms of depth and structure.

As such, results from a CIBIC-plus reflect clinical experience from the trial or trials in which it was used and cannot be compared directly with the results of CIBIC-plus evaluations from other clinical trials. The CIBIC-plus used in Donepezil Hydrochloride Tablets trials was a semi-structured instrument that was intended to examine four major areas of patient function: General, Cognitive, Behavioral and Activities of Daily Living. It represents the assessment of a skilled clinician based upon his/her observations at an interview with the patient, in combination with information supplied by a caregiver familiar with the behavior of the patient over the interval rated. The CIBIC-plus is scored as a seven point categorical rating, ranging from a score of 1, indicating "markedly improved," to a score of 4, indicating "no change" to a score of 7, indicating "markedly worse." The CIBIC-plus has not been systematically compared directly to assessments not using information from caregivers (CIBIC) or other global methods.

### Thirty-Week Study

In a study of 30 weeks duration, 473 patients were randomized to receive single daily doses of placebo, 5 mg/day or 10 mg/day of Donepezil Hydrochloride Tablets. The 30-week study was divided into a 24-week double-blind active treatment phase followed by a 6-week single-blind placebo washout period. The study was designed to compare 5 mg/day or 10 mg/day fixed doses of Donepezil Hydrochloride Tablets to placebo. However, to reduce the likelihood of cholinergic effects, the 10 mg/day treatment was started following an initial 7-day treatment with 5 mg/day doses.

### *Effects on the ADAS-cog*

Figure 1 illustrates the time course for the change from baseline in ADAS-cog scores for all three dose groups over the 30 weeks of the study. After 24 weeks of treatment, the mean differences in the ADAS-cog change scores for Donepezil Hydrochloride Tablets treated patients compared to the patients on placebo were 2.8 and 3.1 points for the

5 mg/day and 10 mg/day treatments, respectively. These differences were statistically significant. While the treatment effect size may appear to be slightly greater for the 10 mg/day treatment, there was no statistically significant difference between the two active treatments.

Following 6 weeks of placebo washout, scores on the ADAS-cog for both the Donepezil Hydrochloride Tablets treatment groups were indistinguishable from those patients who had received only placebo for 30 weeks. This suggests that the beneficial effects of Donepezil Hydrochloride Tablets abate over 6 weeks following discontinuation of treatment and do not represent a change in the underlying disease. There was no evidence of a rebound effect 6 weeks after abrupt discontinuation of therapy.

Figure 1. Time-course of the Change from Baseline in ADAS-cog Score for Patients Completing 24 Weeks of Treatment.





Figure 2 illustrates the cumulative percentages of patients from each of the three treatment groups who had attained the measure of improvement in ADAS-cog score shown on the X axis. Three change scores, (7-point and 4-point reductions from baseline or no change in score) have been identified for illustrative purposes, and the percent of patients in each group achieving that result is shown in the inset table.

The curves demonstrate that both patients assigned to placebo and Donepezil Hydrochloride Tablets have a wide range of responses, but that the active treatment groups are more likely to show greater improvements. A curve for an effective treatment would be shifted to the left of the curve for placebo, while an ineffective or deleterious treatment would be superimposed upon or shifted to the right of the curve

for placebo.


Figure 2. Cumulative Percentage of Patients Completing 24 Weeks of Double-blind Treatment with Specified Changes from Baseline ADAS-cog Scores. The Percentages of Randomized Patients who Completed the Study were: Placebo 80%, 5 mg/day 85% and 10 mg/day 68%.



### *Effects on the CIBIC-plus*

Figure 3 is a histogram of the frequency distribution of CIBIC-plus scores attained by patients assigned to each of the three treatment groups who completed 24 weeks of treatment. The mean drug-placebo differences for these groups of patients were 0.35 points and 0.39 points for 5 mg/day and 10 mg/day of Donepezil Hydrochloride Tablets, respectively. These differences were statistically significant. There was no statistically significant difference between the two active treatments.

Figure 3. Frequency Distribution of CIBIC plus Scores at Week 24.




### Fifteen-Week Study

In a study of 15 weeks duration, patients were randomized to receive single daily doses of placebo or either 5 mg/day or 10 mg/day of Donepezil Hydrochloride Tablets for 12 weeks, followed by a 3-week placebo washout period. As in the 30-week study, to avoid acute cholinergic effects, the 10 mg/day treatment followed an initial 7-day treatment with 5 mg/day doses.

### *Effects on the ADAS-Cog*

Figure 4 illustrates the time course of the change from baseline in ADAS-cog scores for all three dose groups over the 15 weeks of the study. After 12 weeks of treatment, the differences in mean ADAS-cog change scores for the Donepezil Hydrochloride Tablets treated patients compared to the patients on placebo were 2.7 and 3.0 points each, for the 5 and 10 mg/day Donepezil Hydrochloride Tablets treatment groups, respectively. These differences were statistically significant. The effect size for the 10 mg/day group may appear to be slightly larger than that for 5 mg/day. However, the differences between active treatments were not statistically significant.

Figure 4. Time-course of the Change from Baseline in ADAS-cog Score for Patients Completing the 15-week Study.




Following 3 weeks of placebo washout, scores on the ADAS-cog for both the Donepezil Hydrochloride Tablets treatment groups increased, indicating that discontinuation of Donepezil Hydrochloride Tablets resulted in a loss of its treatment effect. The duration of this placebo washout period was not sufficient to characterize the rate of loss of the treatment effect, but, the 30-week study (see above) demonstrated that treatment effects associated with the use of Donepezil Hydrochloride Tablets abate within 6 weeks of treatment discontinuation.

Figure 5 illustrates the cumulative percentages of patients from each of the three treatment groups who attained the measure of improvement in ADAS-cog score shown on the X axis. The same three change scores, (7-point and 4-point reductions from baseline or no change in score) as selected for the 30-week study have been used for this illustration. The percentages of patients achieving those results are shown in the inset table.

As observed in the 30-week study, the curves demonstrate that patients assigned to either placebo or to Donepezil Hydrochloride Tablets have a wide range of responses, but that the Donepezil Hydrochloride Tablets treated patients are more likely to show greater improvements in cognitive performance.


Figure 5. Cumulative Percentage of Patients with Specified Changes from Baseline ADAS-cog Scores. The Percentages of Randomized Patients within Each Treatment Group Who Completed the Study Were: Placebo 93%, 5 mg/day 90% and 10 mg/day 82%.



### Effects on the CIBIC-plus

Figure 6 is a histogram of the frequency distribution of CIBIC-plus scores attained by patients assigned to each of the three treatment groups who completed 12 weeks of treatment. The differences in mean scores for Donepezil Hydrochloride Tablets treated patients compared to the patients on placebo at Week 12 were 0.36 and 0.38 points for the 5 mg/day and 10 mg/day treatment groups, respectively. These differences were statistically significant.

Figure 6. Frequency Distribution of CIBIC plus Scores at Week 12.



In both studies, patient age, sex and race were not found to predict the clinical outcome of Donepezil Hydrochloride Tablets treatment.

## **14.2 Moderate to Severe Alzheimer's Disease**

The effectiveness of Donepezil Hydrochloride Tablets in the treatment of patients with moderate to severe Alzheimer's disease was established in studies employing doses of 10 mg/day and 23 mg/day. Results of a controlled clinical trial in moderate to severe Alzheimer's Disease that compared Donepezil Hydrochloride Tablets 23 mg once daily to 10 mg once daily suggest that a 23 mg dose of Donepezil Hydrochloride Tablets provided additional benefit.

### **Swedish 6 Month Study (10 mg/day)**

The effectiveness of Donepezil Hydrochloride Tablets as a treatment for severe Alzheimer's disease is demonstrated by the results of a randomized, double-blind, placebo-controlled clinical study conducted in Sweden (6 month study) in patients with probable or possible Alzheimer's disease diagnosed by NINCDS-ADRDA and DSM-IV criteria, MMSE: range of 1-10. Two hundred and forty eight (248) patients with severe Alzheimer's disease were randomized to Donepezil Hydrochloride Tablets or placebo. For patients randomized to Donepezil Hydrochloride Tablets, treatment was initiated at 5 mg once daily for 28 days and then increased to 10 mg once daily. At the end of the 6 month treatment period, 90.5% of the Donepezil Hydrochloride Tablets treated patients were receiving the 10 mg/day dose. The mean age of patients was 84.9 years, with a range of 59 to 99. Approximately 77% of patients were women, and 23% were men. Almost all patients were Caucasian. Probable AD was diagnosed in the majority of the patients (83.6% of Donepezil Hydrochloride Tablets treated patients and 84.2% of placebo treated patients).

#### Study Outcome Measures

The effectiveness of treatment with Donepezil Hydrochloride Tablets was determined using a dual outcome assessment strategy that evaluated cognitive function using an instrument designed for more impaired patients and overall function through caregiver-rated assessment. This study showed that patients on Donepezil Hydrochloride Tablets experienced significant improvement on both measures compared to placebo.

The ability of Donepezil Hydrochloride Tablets to improve cognitive performance was assessed with the Severe Impairment Battery (SIB). The SIB, a multi-item instrument, has been validated for the evaluation of cognitive function in patients with moderate to severe dementia. The SIB evaluates selective aspects of cognitive performance, including elements of memory, language, orientation, attention, praxis, visuospatial ability, construction, and social interaction. The SIB scoring range is from 0 to 100, with lower scores indicating greater cognitive impairment.

Daily function was assessed using the Modified Alzheimer's Disease Cooperative Study Activities of Daily Living Inventory for Severe Alzheimer's Disease (ADCS-ADL-severe). The ADCS-ADL-severe is derived from the Alzheimer's Disease Cooperative Study Activities of Daily Living Inventory, which is a comprehensive battery of ADL questions



used to measure the functional capabilities of patients. Each ADL item is rated from the highest level of independent performance to complete loss. The ADCS-ADL-severe is a subset of 19 items, including ratings of the patient's ability to eat, dress, bathe, use the telephone, get around (or travel), and perform other activities of daily living; it has been validated for the assessment of patients with moderate to severe dementia. The ADCS-ADL-severe has a scoring range of 0 to 54, with the lower scores indicating greater functional impairment. The investigator performs the inventory by interviewing a caregiver, in this study a nurse staff member, familiar with the functioning of the patient.

### *Effects on the SIB*

Figure 7 shows the time course for the change from baseline in SIB score for the two treatment groups over the 6 months of the study. At 6 months of treatment, the mean difference in the SIB change scores for Donepezil Hydrochloride Tablets treated patients compared to patients on placebo was 5.9 points. Donepezil Hydrochloride Tablets treatment was statistically significantly superior to placebo.

Figure 7. Time Course of the Change from Baseline in SIB Score for Patients Completing 6 months of Treatment.




Figure 8 illustrates the cumulative percentages of patients from each of the two treatment groups who attained the measure of improvement in SIB score shown on the X-axis. While patients assigned both to Donepezil Hydrochloride Tablets and to placebo have a wide range of responses, the curves show that the Donepezil Hydrochloride

Tablets group is more likely to show a greater improvement in cognitive performance.

Figure 8. Cumulative Percentage of Patients Completing 6 Months of Double-blind Treatment with Particular Changes from Baseline in SIB Scores.





Figure 9. Time Course of the Change from Baseline in ADCS-ADL-Severe Score for Patients Completing 6 Months of Treatment.




#### *Effects on the ADCS-ADL-severe*

Figure 9 illustrates the time course for the change from baseline in ADCS-ADL-severe scores for patients in the two treatment groups over the 6 months of the study. After 6 months of treatment, the mean difference in the ADCS-ADL-severe change scores for Donepezil Hydrochloride Tablets treated patients compared to patients on placebo was 1.8 points. Donepezil Hydrochloride Tablets treatment was statistically significantly superior to placebo.

Figure 10 shows the cumulative percentages of patients from each treatment group with specified changes from baseline ADCS-ADL-severe scores. While both patients assigned to Donepezil Hydrochloride Tablets and placebo have a wide range of responses, the curves demonstrate that the Donepezil Hydrochloride Tablets group is more likely to show a smaller decline or an improvement.

Figure 10. Cumulative Percentage of Patients Completing 6 Months of Double-blind Treatment with Particular Changes from Baseline in ADCS-ADL-Severe Scores.



### Japanese 24-Week Study (10 mg/day)

In a study of 24 weeks duration conducted in Japan, 325 patients with severe Alzheimer's disease were randomized to doses of 5 mg/day or 10 mg/day of donepezil, administered once daily, or placebo. Patients randomized to treatment with donepezil were to achieve their assigned doses by titration, beginning at 3 mg/day, and extending over a maximum of 6 weeks. Two hundred and forty eight (248) patients completed the study, with similar proportions of patients completing the study in each treatment group. The primary efficacy measures for this study were the SIB and CIBIC-plus. At 24 weeks of treatment, statistically significant treatment differences were observed between the 10 mg/day dose of donepezil and placebo on both the SIB and CIBIC-plus. The 5 mg/day dose of donepezil showed a statistically significant superiority to placebo on the SIB, but not on the CIBIC-plus.

### Study of 23 mg/day

The effectiveness of Donepezil Hydrochloride Tablets 23 mg/day as a treatment for moderate to severe Alzheimer's disease has been demonstrated by the results of a randomized, double-blind, controlled clinical investigation in patients with moderate to severe Alzheimer's disease. The controlled clinical study was conducted globally in patients with probable Alzheimer's disease diagnosed by NINCDS-ADRDA and DSM-IV criteria, MMSE: range of 0-20. Patients were required to have been on a stable dose of Donepezil Hydrochloride Tablets 10 mg/day for at least 3 months prior to screening. One thousand four hundred and thirty four (1,434) patients with moderate to severe Alzheimer's disease were randomized to 23 mg/day or 10 mg/day. The mean age of patients was 73.8 years, with a range of 47 to 90. Approximately 63% of patients were

women, and 37% were men. Approximately 36% of the patients were taking memantine throughout the study.

### Study Outcome Measures

The effectiveness of treatment with 23 mg/day was determined using a dual outcome assessment strategy that evaluated cognitive function using an instrument designed for more impaired patients and overall function through caregiver-rated assessment.

The ability of 23 mg/day to improve cognitive performance was assessed with the Severe Impairment Battery (SIB). The SIB, a multi-item instrument, has been validated for the evaluation of cognitive function in patients with moderate to severe dementia. The SIB evaluates selective aspects of cognitive performance, including elements of memory, language, orientation, attention, praxis, visuospatial ability, construction, and social interaction. The SIB scoring range is from 0 to 100, with lower scores indicating greater cognitive impairment.

The ability of 23 mg/day to produce an overall clinical effect was assessed using a Clinician's Interview-Based Impression of Change that incorporated the use of caregiver information, the CIBIC-plus. The CIBIC-plus used in this trial was a semi-structured instrument that examines four major areas of patient function: General, Cognitive, Behavioral and Activities of Daily Living. It represents the assessment of a skilled clinician based upon his/her observations at an interview with the patient, in combination with information supplied by a caregiver familiar with the behavior of the patient over the interval rated. The CIBIC-plus is scored as a seven-point categorical rating, ranging from a score of 1, indicating "markedly improved," to a score of 4, indicating "no change" to a score of 7, indicating "markedly worse."

### *Effects on the SIB*

Figure 11 shows the time course for the change from baseline in SIB score for the two treatment groups over the 24 weeks of the study. At 24 weeks of treatment, the LS mean difference in the SIB change scores for 23 mg/day-treated patients compared to patients treated with 10 mg was 2.2 units ( $p = 0.0001$ ). The dose of 23 mg/day was statistically significantly superior to the dose of 10 mg/day.

Figure 11. Time-course of the Change from Baseline in SIB Score for Patients Completing 24 Weeks of Treatment.





Figure 12 illustrates the cumulative percentages of patients from each of the two treatment groups who attained the measure of improvement in SIB score shown on the X-axis. While patients assigned both to 23 mg/day and to 10 mg/day have a wide range of responses, the curves show that the 23 mg-group is more likely to show a greater improvement in cognitive performance. When such curves are shifted to the left, this indicates a greater percentage of patients responding to treatment on the SIB.


Figure 12. Cumulative Percentage of Patients Completing 24 Weeks of Double-blind Treatment with Specified Changes from Baseline SIB Scores.



*Effects on the CIBIC-plus*

Figure 13 is a histogram of the frequency distribution of CIBIC-plus scores attained by patients at the end of 24 weeks of treatment. The mean difference between the 23 mg/day and 10 mg/day treatment groups was 0.06 units. This difference was not statistically significant.

Figure 13. Frequency Distribution of CIBIC plus Scores at Week 24.



## 16 HOW SUPPLIED/STORAGE AND HANDLING

### 16.1 Donepezil Hydrochloride Tablets

Supplied as film-coated, round tablets containing 5 mg or 10 mg of donepezil hydrochloride.

The 5 mg tablets are white, round, biconvex film coated tablets, debossed 'C57' on one side and plain on other side.

Bottles of 30 (NDC# 71209-019-01)

Bottles of 90 (NDC# 71209-019-04)

Bottles of 1,000 (NDC# 71209-019-11)

The 10 mg tablets are yellow to light yellow, round, biconvex film coated tablets, debossed 'C55' on one side and plain on other side.

Bottles of 30 (NDC# 71209-020-01)

Bottles of 90 (NDC# 71209-020-04)

Bottles of 1,000 (NDC# 71209-020-11)

#### Storage

Store at controlled room temperature, 15°C to 30°C (59°F to 86°F).

## 17 PATIENT COUNSELING INFORMATION



Advise the patient to read the FDA-approved patient labeling (Patient Information). Instruct patients and caregivers to take Donepezil Hydrochloride Tablets only once per day, as prescribed. Instruct patients and caregivers that Donepezil Hydrochloride Tablets can be taken with or without food. Advise patients and caregivers that Donepezil Hydrochloride Tablets may cause nausea, diarrhea, insomnia, vomiting, muscle cramps, fatigue, and decreased appetite. Advise patients to notify their healthcare provider if they are pregnant or plan to become pregnant.

**Manufactured By:**  
**Cadila Pharmaceuticals Limited**  
1389, Trasad Road,  
Dholka - 382225,  
District - Ahmedabad, Gujarat, INDIA

Revised: November, 2019

## **DONEPEZIL HYDROCHLORIDE TABLETS PATIENT PACKAGE INSERT**

Donepezil Hydrochloride (doe-NEP-e-zil HYE-droe-KLOR-ide) Tablets

- Tablets: 5 mg and 10 mg

Read this Patient Information that comes with Donepezil Hydrochloride Tablets before you start taking it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with the doctor about Alzheimer's disease or treatment for it. If you have questions, ask the doctor or pharmacist.

### **What is Donepezil Hydrochloride Tablets?**

Donepezil Hydrochloride Tablets comes as Donepezil Hydrochloride film-coated tablets in dosage strengths of 5 mg and 10 mg.

Donepezil Hydrochloride Tablets is a prescription medicine to treat mild, moderate and severe Alzheimer's disease. Donepezil Hydrochloride Tablets can help with mental function and with doing daily tasks. Donepezil Hydrochloride Tablets does not work the same in all people. Some people may:

- Seem much better
- Get better in small ways or stay the same
- Get worse over time but slower than expected
- Not change and then get worse as expected

Donepezil Hydrochloride does not cure Alzheimer's disease. All patients with Alzheimer's

disease get worse over time, even if they take Donepezil Hydrochloride. Donepezil Hydrochloride has not been approved as a treatment for any medical condition in children.

### **Who should not take Donepezil Hydrochloride Tablets?**

Do not take Donepezil Hydrochloride Tablets if you are allergic to any of the ingredients in Donepezil Hydrochloride Tablets or to medicines that contain piperidines. Ask your doctor if you are not sure. See the end of this leaflet for a list of ingredients in Donepezil Hydrochloride Tablets.

### **What should I tell my doctor before taking Donepezil Hydrochloride Tablets?**

**Tell the doctor about all of your present or past health problems and conditions.** Include:

- Any heart problems including problems with irregular, slow, or fast heartbeats
- Asthma or lung problems
- A seizure
- Stomach ulcers
- Difficulty passing urine
- Liver or kidney problems
- Trouble swallowing tablets
- Present pregnancy or plans to become pregnant. It is not known if Donepezil Hydrochloride Tablets can harm an unborn baby.
- Present breast-feeding. It is not known if Donepezil Hydrochloride Tablets passes into breast milk. Talk to your doctor about the best way to feed your baby if you take Donepezil Hydrochloride Tablets.

**Tell the doctor about all the medicines you take,** including prescription and non-prescription medicines, vitamins, and herbal products. Donepezil Hydrochloride Tablets and other medicines may affect each other.

Be particularly sure to tell the doctor if you take aspirin or medicines called nonsteroidal anti-inflammatory drugs (NSAIDs). There are many NSAID medicines, both prescription and non-prescription. Ask the doctor or pharmacist if you are not sure if any of your medicines are NSAIDs. Taking NSAIDs and Donepezil Hydrochloride Tablets together may make you more likely to get stomach ulcers.

Donepezil Hydrochloride Tablets taken with certain medicines used for anesthesia may cause side effects. Tell the responsible doctor or dentist that you take Donepezil Hydrochloride Tablets before you have:

- surgery
- medical procedures
- dental surgery or procedures.

Know the medicines that you take. Keep a list of all your medicines. Show it to your or

pharmacist before you start a new medicine.

### **How should you take Donepezil Hydrochloride Tablets?**

- Take Donepezil Hydrochloride Tablets exactly as prescribed by the doctor. Do not stop Donepezil Hydrochloride Tablets or change the dose yourself. Talk with your doctor first.
- Take Donepezil Hydrochloride Tablets one time each day. Donepezil Hydrochloride Tablets can be taken with or without food.
- If you miss a dose of Donepezil Hydrochloride Tablets, just wait. Take only the next dose at the usual time. Do not take 2 doses at the same time.
- If Donepezil Hydrochloride Tablets is missed for 7 days or more, talk with your doctor before starting again.
- If you take too much Donepezil Hydrochloride Tablets at one time, call your doctor or poison control center, or go to the emergency room right away.

### **What are the possible side effects of Donepezil Hydrochloride Tablets?**

#### **Donepezil Hydrochloride Tablets may cause the following serious side effects:**

- slow heartbeat and fainting. This happens more often in people with heart problems. Call your doctor right away if you feel faint or lightheaded while taking Donepezil Hydrochloride Tablets.
- more stomach acid. This raises the chance of ulcers and bleeding. The risk is higher for people who have had ulcers, or take aspirin or other NSAIDs.
- worsening of lung problems in people with asthma or other lung disease.
- seizures.
- difficulty passing urine.

#### **Call your doctor *right away* if you have:**

- fainting.
- heartburn or stomach pain that is new or won't go away.
- nausea or vomiting, blood in the vomit, dark vomit that looks like coffee grounds.
- bowel movements or stools that look like black tar.
- new or worse asthma or breathing problems.
- seizures.
- difficulty passing urine.

#### **The most common side effects of Donepezil Hydrochloride Tablets are:**

- nausea
- diarrhea
- not sleeping well
- vomiting

- muscle cramps
- feeling tired
- not wanting to eat

**These side effects may get better after you take Donepezil Hydrochloride Tablets for a while. This is not a complete list of side effects with Donepezil Hydrochloride Tablets. For more information, ask your doctor or pharmacist.**

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

#### **How should Donepezil Hydrochloride Tablets be stored?**

Store Donepezil Hydrochloride Tablets at room temperature between 59° to 86°F (15° to 30°C).

**Keep Donepezil Hydrochloride Tablets and all medicines out of the reach of children.**

#### **General information about Donepezil Hydrochloride Tablets**

Medicines are sometimes prescribed for conditions that are not mentioned in this Patient Information Leaflet. Do not use Donepezil Hydrochloride Tablets for a condition for which it was not prescribed. Do not give Donepezil Hydrochloride Tablets to other people, even if they have the same symptoms or condition. It may harm them.

This leaflet summarizes the most important information about Donepezil Hydrochloride Tablets. If you would like more information talk with your doctor. You can ask your pharmacist or doctor for information about Donepezil Hydrochloride Tablets that is written for health professionals. For more information, call Cadila Pharmaceuticals Limited at 1-202-355-9784.

#### **What are the ingredients in Donepezil Hydrochloride Tablets?**

**Active ingredient:** donepezil hydrochloride

#### **Inactive ingredients:**

- **Donepezil Hydrochloride Tablets 5 mg and 10 mg film-coated tablets:** lactose monohydrate, pregelatinized starch, microcrystalline cellulose, hydroxypropyl cellulose, and magnesium stearate. The film coating contains talc, polyethylene glycol, hypromellose, and titanium dioxide. Additionally, the 10 mg tablet contains yellow iron oxide as a coloring agent.

#### **Manufactured By:**

**Cadila Pharmaceuticals Limited**

1389, Trasad Road,

Dholka - 382225,

District - Ahmedabad, Gujarat, INDIA

Revised: November, 2019

**Rx Only**

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**


NDC 71209-019-01

**Donepezil Hydrochloride Tablets USP**

**5 mg**

RX Only

30 Tablets



**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**



NDC 71209-019-04

**Donepezil Hydrochloride Tablets USP**

**5 mg**

RX Only

90 Tablets

<p>Each film-coated tablet contains : Donepezil hydrochloride USP 5 mg.</p> <p><b>DOSAGE AND USE</b> : See accompanying prescribing information.</p> <p>Store at controlled room temperature 59° to 86°F (15°C to 30°C).</p> <p>Dispense in tight containers (USP).</p> <p>Manufactured by : <b>CADILA PHARMACEUTICALS LTD.</b> 1389, Dholka, Dist. Ahmedabad, Gujarat State, INDIA.</p>	<p style="text-align: center;">NDC 71209-019-04</p> <p style="text-align: center;"><b>Donepezil Hydrochloride Tablets USP</b></p> <p style="text-align: center;"><b>5 mg</b></p> <p style="text-align: center;">Rx Only</p> <p style="text-align: right;">90 Tablets</p> 	 <p style="text-align: center;">J00000X</p> <p style="text-align: center;"><i>Unvarnished Area 32 x 25 mm Space for serialized datamatrix code and batch coding.</i></p>
--	---	---

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**



NDC 71209-019-11

**Donepezil Hydrochloride Tablets USP**

**5 mg**

RX Only

1000 Tablets

<p>Each film-coated tablet contains : Donepezil hydrochloride USP 5 mg.</p> <p><b>DOSAGE AND USE</b> : See accompanying prescribing information.</p> <p>Store at controlled room temperature 59° to 86°F (15°C to 30°C).</p> <p>Dispense in tight containers (USP).</p> <p>Manufactured by : <b>CADILA PHARMACEUTICALS LTD.</b> 1389, Dholka, Dist. Ahmedabad, Gujarat State, INDIA.</p>	<p style="text-align: center;">NDC 71209-019-11</p> <p style="text-align: center;"><b>Donepezil Hydrochloride Tablets USP</b></p> <p style="text-align: center;"><b>5 mg</b></p> <p style="text-align: center;">Rx Only</p> <p style="text-align: right;">1000 Tablets</p> 	 <p style="text-align: center;">J00000X</p> <p style="text-align: center;"><i>Unvarnished Area 65 x 25 mm Space for serialized datamatrix code and batch coding.</i></p>
--	---	---

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 71209-020-01

**Donepezil Hydrochloride Tablets USP**

**10 mg**

RX Only

30 Tablets

<p>Each film-coated tablet contains : Donepezil hydrochloride USP 10 mg.</p> <p><b>DOSAGE AND USE :</b> See accompanying prescribing information.</p> <p>Store at controlled room temperature 59° to 86°F (15°C to 30°C).</p> <p>Dispense in tight containers (USP).</p> <p>Manufactured by : <b>CADILA PHARMACEUTICALS LTD.</b> 1389, Dholka, Dist. Ahmedabad, Gujarat State, INDIA.</p>	<p style="text-align: center;">NDC 71209-020-01</p> <p style="text-align: center;"><b>Donepezil Hydrochloride Tablets USP</b></p> <p style="text-align: center;"><b>10 mg</b></p> <p style="text-align: center;">Rx Only</p> <p style="text-align: right;">30 Tablets</p> 		<p style="color: red; text-align: center;"><i>Unvarnished Area 32 x 25 mm Space for serialized datamatrix code and batch coding.</i></p>
---	--	---	--

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**


NDC 71209-020-04

**Donepezil Hydrochloride Tablets USP**

**10 mg**

RX Only

90 Tablets

<p>Each film-coated tablet contains : Donepezil hydrochloride USP 10 mg.</p> <p><b>DOSAGE AND USE :</b> See accompanying prescribing information.</p> <p>Store at controlled room temperature 59° to 86°F (15°C to 30°C).</p> <p>Dispense in tight containers (USP).</p> <p>Manufactured by : <b>CADILA PHARMACEUTICALS LTD.</b> 1389, Dholka, Dist. Ahmedabad, Gujarat State, INDIA.</p>	<p style="text-align: center;">NDC 71209-020-04</p> <p style="text-align: center;"><b>Donepezil Hydrochloride Tablets USP</b></p> <p style="text-align: center;"><b>10 mg</b></p> <p style="text-align: center;">Rx Only</p> <p style="text-align: right;">90 Tablets</p> 		<p style="color: red; text-align: center;"><i>Unvarnished Area 32 x 25 mm Space for serialized datamatrix code and batch coding.</i></p>
---	--	---	--

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**



NDC 71209-020-11

# Donepezil Hydrochloride Tablets USP

10 mg

RX Only

1000 Tablets

<p>Each film-coated tablet contains : Donepezil hydrochloride USP 10 mg.</p> <p><b>DOSAGE AND USE</b> : See accompanying prescribing information.</p> <p>Store at controlled room temperature 59° to 86°F (15°C to 30°C).</p> <p>Dispense in tight containers (USP).</p> <p>Manufactured by : <b>CADILA PHARMACEUTICALS LTD.</b> 1389, Dholka, Dist. Ahmedabad, Gujarat State, INDIA.</p>	<p>NDC 71209-020-11</p> <p><b>Donepezil Hydrochloride Tablets USP</b></p> <p><b>10 mg</b></p> <p><b>Rx Only</b></p> <p><b>1000 Tablets</b></p> 	 <p>31171209020114</p> <p>000000</p>	<p>Unvarnished Area 65 x 25 mm Space for serialized datamatrix code and batch coding.</p>
---	--	--	---

## DONEPEZIL HYDROCHLORIDE

donepezil hydrochloride tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:71209-019
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DONEPEZIL HYDROCHLORIDE</b> (UNII: 3O2T2PJ89D) (DONEPEZIL - UNII:8SSC91326P)	DONEPEZIL HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>HYDROXYPROPYL CELLULOSE</b> (UNII: RFW2ET671P)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)	



**TITANIUM DIOXIDE** (UNII: 15FIX9V2JP)

### Product Characteristics

<b>Color</b>	WHITE (WHITE)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND,BICONVEX)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	C57
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71209-019-01	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2018	
2	NDC:71209-019-04	90 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2018	
3	NDC:71209-019-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204609	01/30/2017	

## DONEPEZIL HYDROCHLORIDE

donepezil hydrochloride tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:71209-020
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DONEPEZIL HYDROCHLORIDE</b> (UNII: 3O2T2PJ89D) (DONEPEZIL - UNII:8SSC91326P)	DONEPEZIL HYDROCHLORIDE	10 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>HYDROXYPROPYL CELLULOSE</b> (UNII: RFW2ET671P)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	

<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>HYPROMELLOSE 2910 (6 MPA.S)</b> (UNII: 0WZ8WG20P6)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)	

### Product Characteristics

<b>Color</b>	YELLOW (YELLOW TO LIGHT YELLOW)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND, BICONVEX)	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	C55
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71209-020-01	30 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2018	
2	NDC:71209-020-04	90 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2018	
3	NDC:71209-020-11	1000 in 1 BOTTLE; Type 0: Not a Combination Product	11/24/2018	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204609	01/30/2017	

**Labeler** - Cadila Pharmaceuticals Limited (862257719)

### Establishment

Name	Address	ID/FEI	Business Operations
Cadila Pharmaceuticals Limited		918451696	ANALYSIS(71209-019, 71209-020) , LABEL(71209-019, 71209-020) , MANUFACTURE(71209-019, 71209-020) , PACK(71209-019, 71209-020)

Revised: 12/2022

Cadila Pharmaceuticals Limited