

**KAOPECTATE- bismuth subsalicylate tablet, coated**  
**Kramer Laboratories**

-----  
**Drug Facts**

<b>Active ingredient (per caplet)</b>	<b>Purposes</b>
Bismuth subsalicylate 262 mg	Anti-diarrheal, Upset stomach reliever

**Uses** relieves:

- traveler's diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - nausea
  - gas
  - belching
  - fullness

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert: Contains salicylate.**

**Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

**Ask a doctor or pharmacist before use if you are** taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### ***Directions***

- **adults and children 12 years of age and older:** 2 caplets (1 dose)
- swallow caplets with water; do not chew
- repeat dose every 1/2 hour to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### ***Other information***

- Each caplet contains very low sodium
- **do not use if carton was opened or if printed foil inner seal under bottle cap is torn or missing**
- store at room temperature 20°-25° C (68°-77° F)

### ***Inactive ingredients***

croscarmellose sodium, crospovidone, hydrated silica, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, povidone, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin

### **Principal Display Panel**

***FAST & EFFECTIVE RELIEF***

**Kaopectate®**

*Bismuth Subsalicylate*

• *Anti-Diarrheal* • *Upset Stomach Reliever*

## **Diarrhea & Upset Stomach**

- ✓ Begins controlling symptoms from the first dose
- ✓ Quickly relieves urgency, gas, and cramping
- ✓ Effective on diarrhea from bacteria, viruses, and other causes

CLINICALLY PROVEN MEDICINE

28 Coated Caplets



**LOT**

**EXP**

**Do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken.**

**Kramer Laboratories, Inc.**

Bridgewater, NJ 08807

**1-800-824-4894**

14711-07-20\_K0720

# Kaopectate®

**Bismuth Subsalicylate**  
• Anti-Diarrheal • Upset Stomach Reliever

## Drug Facts

**Active ingredient (per caplet)** Bismuth subsalicylate 262 mg.....**Purposes** Anti-diarrheal, Upset stomach reliever

**Uses** relieves: • traveler's diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy Alert:** Contains salicylate. Do not take if you are  
• allergic to salicylates (including aspirin) • taking other salicylate products

## Drug Facts (continued)

**Do not use if you have** • an ulcer • a bleeding problem • bloody or black stool

**Ask a doctor before use if you have** • fever • mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for**  
• anticoagulation (thinning the blood) • diabetes • gout • arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if** • symptoms get worse  
• ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## Drug Facts (continued)

### Directions

- adults and children 12 years of age and older: 2 caplets (1 dose)
- swallow caplets with water; do not chew
- repeat dose every 1/2 hour to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### Other information

- Each caplet contains very low sodium
- do not use if carton was opened or if printed foil inner seal under bottle cap is torn or missing
- store at room temperature 20°-25° C (68°-77° F)

## Drug Facts (continued)

### Inactive ingredients


croscarmellose sodium, crospovidone, hydrated silica, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, povidone, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin

**Kramer Laboratories, Inc.**  
Bridgewater, NJ 08807  
1-800-824-4894

14711-07-20\_K0720

Do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken.

LOT  
EXP



**Kaopectate®**

**Bismuth Subsalicylate**

- **Anti-Diarrheal**
- **Upset Stomach Reliever**

**Diarrhea & Upset Stomach**

**28 Coated Caplets**

**Do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken.**

**KEEP CARTON AS IT CONTAINS IMPORTANT INFORMATION**

**Kramer Laboratories, Inc.**

Bridgewater, NJ 08807

**1-800-824-4894**

14712-07-20\_K0720



**Active ingredient** (in each caplet)  
Bismuth subsalicylate 262 mg

**Purposes** Anti-diarrheal, Upset stomach  
reliever **Uses relieves:** • traveler's  
diarrhea • diarrhea • upset stomach  
due to overindulgence in food and drink,  
including: • heartburn • indigestion •  
nausea • gas • belching • fullness

**Warnings** **Reye's syndrome:** Children  
and teenagers who have or are recovering  
from chicken pox or flu-like symptoms  
should not use this product. When using  
this product, if changes in behavior with  
nausea and vomiting occur, consult a  
doctor because these symptoms could  
be an early sign of Reye's syndrome, a  
rare but serious illness. **Allergy alert:**  
**Contains salicylate. Do not take if you**  
**are** • allergic to salicylates (including  
aspirin) • taking other salicylate products  
**Do not use if you have** • an  
ulcer • a bleeding problem  
• bloody or black stool **Ask**  
**a doctor before use if you**  
**have** • fever • mucus in the stool **Ask a**  
**doctor or pharmacist before use if you**  
**are taking any drug for** • anticoagulation  
(thinning the blood) • diabetes • gout •

**PEEL HERE ▼**

arthritis

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** • adults and children 12 years of age and older: 2 caplets (1 dose)

- swallow caplets with water; do not chew
- repeat dose every 1/2 hour to 1 hour as needed
- do not exceed 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** • Each caplet contains

Continued ▼

very low sodium • **do not use if carton was opened or if printed foil inner seal under bottle cap is torn or missing**

- store at room temperature 20°-25° C (68°-77° F)

**Inactive ingredients** croscarmellose sodium, crospovidone, hydrated silica, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, povidone, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin

**Do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken.**

**KEEP CARTON AS IT CONTAINS IMPORTANT INFORMATION**

Kramer Laboratories, Inc.  
Bridgewater, NJ 08807  
1-800-824-4894

14712-07-20\_K0720

# KAOPECTATE

bismuth subsalicylate tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55505-197
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Bismuth Subsalsicylate</b> (UNII: 62TEY51RR1) (Bismuth Cation - UNII:ZS9CD1I8YE)	Bismuth Subsalsicylate	262 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>Croscarmellose Sodium</b> (UNII: M28OL1HH48)	
<b>Crospovidone</b> (UNII: 2S7830E561)	
<b>Hydrated Silica</b> (UNII: Y6O7T4G8P9)	
<b>Hypromellose, Unspecified</b> (UNII: 3NXW29V3WO)	
<b>Magnesium Stearate</b> (UNII: 70097M6I30)	
<b>Microcrystalline Cellulose</b> (UNII: OP1R32D61U)	
<b>Mineral Oil</b> (UNII: T5L8T28FGP)	
<b>Povidone, Unspecified</b> (UNII: FZ989GH94E)	
<b>Sodium Lauryl Sulfate</b> (UNII: 368GB5141J)	
<b>Stearic Acid</b> (UNII: 4ELV7Z65AP)	
<b>Talc</b> (UNII: 7SEV7J4R1U)	
<b>Titanium Dioxide</b> (UNII: 15FIX9V2JP)	
<b>Triacetin</b> (UNII: XHX3C3X673)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL (caplet)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	K
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:55505-197-07	1 in 1 CARTON	10/01/2020	
1		28 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55505-197-70	1 in 1 CARTON	02/01/2021	
2		42 in 1 BOTTLE; Type 0: Not a Combination Product		



## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	10/01/2020	

**Labeler** - Kramer Laboratories (122720675)

Revised: 12/2023

Kramer Laboratories