

BISACODYL LAXATIVE - bisacodyl suppository

McKesson

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

ACTIVE INGREDIENT (in each suppository)

Bisacodyl, 10 mg

PURPOSE

Stimulant Laxative

USES

- For relief of occasional constipation
- This product generally produces bowel movement in 15 minutes to 1 hour

WARNINGS

For rectal use only. May cause abdominal discomfort, faintness, rectal burning, and mild cramps.

Do not use

- more than one per day
- for a period of longer than one week unless directed by a doctor
- laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor
- if seal under product lid is torn, damaged, or open

Ask a doctor before use

- if you have noticed a sudden change in bowel habits that persist over a period of two weeks
- if you are pregnant or nursing a baby

Stop use and ask a doctor

if you have rectal bleeding or fail to have a bowel movement after using a laxative. This may indicate a serious condition.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

Adults and children 12 years of age and older	One suppository once daily
Children 6 to 12 years	1/2 suppository once daily
Children under 6	Ask a doctor

- Detach one suppository from the strip and remove from foil
- Carefully insert one suppository well into the rectum
- Do not use more than once per day

OTHER INFORMATION

- Store at room temperature: 15° - 30° C (59° - 86° F). Do not exceed 30° C (86° F).
- Individually sealed for your protection. Do not use if foil is torn or open.

INACTIVE INGREDIENT

hydrogenated vegetable oil

PACKAGE INFORMATION

Sunmark®

COMPARE TO DULCOLAX® ACTIVE INGREDIENT
NDC 49348-598-67

For prompt relief of constipation often in 15 minutes to 1 hour

Laxative Suppositories

BISACODYL USP
CONSTIPATION RELIEF

8 SUPPOSITORIES 10 mg EACH

FOR YOUR SAFETY, SUPPOSITORIES ARE PACKAGED IN PAMPER-EVIDENT SEALED PLASTIC WRAPPER. DO NOT USE IF PLASTIC WRAPPER IS TORN OR OPEN.

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., owner of the registered trademark Dulcolax®.

Another Quality Product
Distributed by McKesson
One Post Street, San Francisco, CA 94104
Money Back Guarantee
Please visit us at www.sunmarkbrand.com

sunmark[®]

COMPARE TO DUCOLAX[®]
ACTIVE INGREDIENT[™]
NDC 49344-398-87

For prompt relief of constipation
often in 15 minutes to 1 hour

sunmark[®]

laxative suppositories

BISACODYL USP

CONSTIPATION RELIEF

8 SUPPOSITORIES 10 mg EACH



MCKESSON
A Division of McKesson
Distribution, Inc.
2000 North Shoreline Blvd., Emeryville, CA 94608
© 2014 McKesson
Please visit us at www.sunmark.com

Rev. A

Drug Facts

Active Ingredient (in each suppository) Bisacodyl 10 mg
Purpose Stimulant Laxative

Uses
 ■ For relief of occasional constipation
 ■ This product generally produces bowel movement in 15 minutes to 1 hour

Warnings
 ■ Do not use more than one per day
 ■ For a period of longer than one week unless directed by a doctor
 ■ Laxative products when absorbed pass, nausea, or vomiting are present unless directed by a doctor
 ■ Avoid use around the suppository if torn, damaged, or open
 ■ Ask a doctor ■ Before use if you have noticed a sudden change in bowel habits that persists over a period of two weeks
 ■ If you are pregnant or nursing a baby
 ■ Stop use and ask a doctor if you have rectal bleeding or fail to have a bowel movement after using a laxative. This may indicate a serious condition.
 ■ Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
 Adults and children 12 years of age and older
 One suppository once daily
 Children 6 to 12 years
 ½ suppository once daily
 Children under 6
 Ask a doctor.

Other Information
 ■ Do not use more than once per day
 ■ Carefully insert one suppository well into the rectum
 ■ Detach one suppository from the strip and remove from the foil wrapper
 ■ Store at room temperature, 15°–30° C (59°–86° F). Do not exceed 30° C (86° F) and avoid freezing for your protection. Do not use if foil is torn or open.

Inactive ingredient hydrogenated vegetable oil

SUPPOSITORIES ARE PACKAGED IN TAMPER-EVIDENT SEALED PLASTIC WRAPPERS. DO NOT USE IF PLASTIC WRAPPERS IS TORN OR OPEN.

By Andrew's Healthcare Therapeutics, Inc.
New York, NY
© 2014

BISACODYL LAXATIVE

bisacodyl suppository

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49348-598
Route of Administration	RECTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
bisacodyl (UNII: 10X0709Y6I) (bisacodyl - UNII:10X0709Y6I)	bisacodyl	10 mg

Inactive Ingredients

Ingredient Name	Strength
fat, hard (UNII: 8334LX7S21)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-598-67	8 in 1 BOX		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part334	08/08/2005	

Labeler - McKesson (177667227)