

**ANTI-DIARRHEAL- loperamide hcl tablet
DOLGENCORP, LLC**

Dollar General 44-375

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use

if you have bloody or black stool.

Ask a doctor before use if you have

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product,

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

1-888-309-9030

Principal Display Panel

DG® | health

Compare to active ingredient of Imodium® A-D*

Anti-Diarrheal
Loperamide HCl Tablets, 2 mg
Anti-Diarrheal

- **Controls the symptoms of diarrhea**

2
mg each

24 Caplets
Actual Caplet Size

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING


1. To open, tear at perforations and remove a section.

2. Use ✂ to cut through plastic for caplet.

*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Imodium® A-D.
50844 ORG061937508

DISTRIBUTED BY
OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

100% Satisfaction
Guaranteed!
(888)309-9030



B-0315-375-08
0R0061937508

Drug Facts (continued)

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use if you have bloody or black stool.

Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea. ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool but no more than 4 caplets in 24 hours
children 9-11 years (50-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

■ **TAMPER EVIDENCE: DO NOT USE IF OUTER PACKAGE IS OPENED OR BUSTER IS TORN OR BROKEN**

■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number

Inactive ingredients corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments? 1-888-309-9030

Drug Facts

Active ingredient (in each caplet) Loperamide HCl 2 mg. Anti-diarrheal

Purpose Loperamide HCl 2 mg. Anti-diarrheal

Use Use to control symptoms of diarrhea, including Traveler's Diarrhea

Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to piperamide HCl.

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION




Compare to active ingredient of Imodium® A-D*

Anti-Diarrheal

Loperamide HCl Tablets, 2 mg

Anti-Diarrheal

- Controls the symptoms of diarrhea




DO NOT USE IF PACKAGE IS TORN, BROKEN OR UNIT IS TORN, BROKEN OR SIGNS OF TAMPERING

No varnish
Exp date

*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Imodium® A-D, 50844 0R0061937508

DISTRIBUTED BY
OLD EAST MAIN CO
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072





ANTI-DIARRHEAL

loperamide hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-375
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X90C3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	44;375
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-375-02	2 in 1 CARTON	01/12/2024	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:55910-375-08	4 in 1 CARTON	01/12/2024	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	01/12/2024	

Labeler - DOLGENCORP, LLC (068331990)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-375) , pack(55910-375)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55910-375)

Revised: 2/2024

DOLGENCORP, LLC