

**MEDS ON THE GO- acetaminophen, calcium carbonate, dextromethorphan hbr, guaifenesin, phenylephrine hcl, ibuprofen, loperamide hcl**

**Doc in the Box LLC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

## **MEDS ON THE GO**

### **EXTRA STRENGTH NON-ASPIRIN**

***Active ingredient (in each tablet)***

Acetaminophen 500 mg

### **EXTRA STRENGTH NON-ASPIRIN**

***Purpose***

Pain reliever/fever reducer

### **EXTRA STRENGTH NON-ASPIRIN**

***Uses***

**For the temporary relief of minor aches and pains associated with**

- headache
- muscular aches
- minor arthritis pain
- common cold
- toothache
- menstrual cramps

**For the reduction of fever.**

### **EXTRA STRENGTH NON-ASPIRIN**

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 8 tablets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **EXTRA STRENGTH NON-ASPIRIN**

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- for more than 3 days for fever unless directed by a doctor

#### **EXTRA STRENGTH NON-ASPIRIN**

**Ask a doctor before use if you have** liver disease

#### **EXTRA STRENGTH NON-ASPIRIN**

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

#### **EXTRA STRENGTH NON-ASPIRIN**

**Stop using and ask a doctor if**

- symptoms do not improve
- new symptoms occur
- pain or fever persists or gets worse
- redness or swelling is present

#### **EXTRA STRENGTH NON-ASPIRIN**

**If pregnant or breast-feeding,** ask a health professional before use.

#### **EXTRA STRENGTH NON-ASPIRIN**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### **EXTRA STRENGTH NON-ASPIRIN**

##### ***Directions***

- **Do not use more than directed**

---

<b>Adults and children: (12 years and older)</b>	Take 2 tablets with water every 4 to 6 hours as needed. Do not take more than 8 tablets in 24 hours.
<b>Children under 12 years:</b>	Do not give this adult strength product to children under 12 years of age; this will provide more than the recommended dose (overdose) and may cause liver damage.

---

#### **EXTRA STRENGTH NON-ASPIRIN**

##### ***Other information***

- store at room temperature 59° - 86°F (15° - 30°C)
- tamper evident sealed packets
- do not use any opened or torn packets

#### **EXTRA STRENGTH NON-ASPIRIN**

##### ***Inactive ingredients***

corn starch, hypromellose, maltodextrin\*, microcrystalline cellulose\*, polyethylene glycol, povidone\*, pregelatinized starch\*, sodium starch glycolate\*, stearic acid, titanium dioxide\*.

\* may contain

## **EXTRA STRENGTH NON-ASPIRIN**

**Questions? 1-800-634-7680**

### **ANTACID**

***Active ingredient (in each tablet)***

Calcium Carbonate 420 mg

### **ANTACID**

***Purpose***

Antacid

### **ANTACID**

***Uses***

For the relief of the following symptoms associated with

- acid indigestion
- sour stomach
- heartburn
- upset stomach

### **ANTACID**

***Warnings***

### **ANTACID**

**Ask a doctor or health professional before use if you have**

- been taking a prescription drug. Antacids may interact with certain prescription drugs
- kidney stones
- a calcium-restricted diet

### **ANTACID**

**Stop using this product and ask a doctor if symptoms last more than 2 weeks**

### **ANTACID**

**Do not exceed recommended dosage.**

**If pregnant or breast-feeding, ask a health professional before use.**

### **ANTACID**

**Keep out of the reach of children.**

## ANTACID

### *Directions*

- **do not use more than directed**
- **Adults and children (12 years and older):** Chew 2 tablets every 2 or 3 hours as symptoms occur or as directed by a physician. Do not take more than 19 tablets in a 24 hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician.
- **Children under 12 years:** Do not give to children under 12 years of age.

## ANTACID

### *Other information*

- Phenylketonurics: contains phenylalanine 1.5 mg per tablet
- each tablet contains 168 mg of elemental calcium
- store at room temperature 59° - 86°F (15° - 30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

## ANTACID

### *Inactive ingredients*

aspartame\*, croscarmellose sodium\*, gum acacia\*, magnesium stearate, maltodextrin, mineral oil\*, mint flavor, sorbitol\*, sucrose\*.

\* may contain

## ANTACID

**Questions or comments? call 1-800-634-7680**

## COLD RELIEF

### *Active ingredient (in each tablet)*

Acetaminophen 325 mg

Dextromethorphan Hydrobromide 15 mg

Guaifenesin 200 mg

Phenylephrine Hydrochloride 5 mg

## COLD RELIEF

### *Purpose*

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

## **COLD RELIEF**

### ***Uses***

#### **Temporarily relieves these cold symptoms**

- cough
- sore throat
- minor aches and pains
- headache
- nasal congestion
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

#### **Temporarily reduces fever.**

## **COLD RELIEF**

### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor promptly.

## **COLD RELIEF**

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## **COLD RELIEF**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough that lasts as occurs with smoking, asthma, chronic bronchitis or emphysema

## COLD RELIEF

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

## COLD RELIEF

**When using this product**

- do not use more than directed

## COLD RELIEF

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain or nasal congestion gets worse or lasts for more than 7 days
- fever gets worse or lasts for more than 3 days
- you get nervous, dizzy or sleepless
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

## COLD RELIEF

**If pregnant or breast-feeding**, ask a health professional before use.

## COLD RELIEF

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## COLD RELIEF

### *Directions*

---

<b>Adults and children: (12 years and older)</b>	Take 2 tablets with water every 6-8 hours as needed. Do not take more than 8 tablets in 24 hours.
<b>Children under 12 years:</b>	Do not give to children under 12 years of age.

---

## COLD RELIEF

### *Other information*

- store at room temperature 59° - 86°F (15° - 30°C)
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

## COLD RELIEF

**Inactive ingredients** maltodextrin, microcrystalline cellulose, povidone, sodium starch glycolate, starch, stearic acid

## **COLD RELIEF**

**Questions?**

**1-800-634-7680**

## **IBUPROFEN**

**Active ingredient (in each tablet)**

Ibuprofen 200 mg (NSAID)\*

\*non-steroidal anti-inflammatory drug

## **IBUPROFEN**

**Purpose**

Pain reliever/fever reducer

## **IBUPROFEN**

**Uses**

**Temporarily relieves minor aches and pains associated with**

- headache
- toothache
- backache
- menstrual cramps
- common cold
- muscular aches
- minor arthritis pain

**Temporarily reduces fever.**

## **IBUPROFEN**

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- skin reddening
- asthma (wheezing)
- facial swelling
- rash
- shock
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- taking a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

## **IBUPROFEN**

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

## **IBUPROFEN**

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

## **IBUPROFEN**

### **Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug
- under a doctor's care for any serious condition

## **IBUPROFEN**

### **When using this product**

- the risk of heart attack or stroke may increase if you use more than directed or longer than directed
- take with food or milk if stomach upset occurs

## **IBUPROFEN**

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
- have stomach pain that does not get better
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts for more than 3 days
  - redness or swelling is present in the painful area
  - any new or unexpected symptoms occur

## **IBUPROFEN**

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to

use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

## **IBUPROFEN**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## **IBUPROFEN**

### ***Directions***

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

---

**Adults and children:  
(12 years and older)** Take 1 tablet every 4 to 6 hours while symptoms persist. If pain for fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

**Children under 12 years:** Do not give to children under 12 years of age.

---

## **IBUPROFEN**

### ***Other information***

- read all product information before using
- store at 68° - 77°F (20° - 25°C)
- avoid excessive heat 104°F (above 40°C)
- tamper evident sealed packets
- do not use any opened or torn packet

## **IBUPROFEN**

### ***Inactive ingredients***

carnauba wax\*, corn starch, hypromellose\*, iron oxide red, lactose\*, magnesium stearate\*, microcrystalline cellulose\*, polydextrose\*, polyethylene glycol, polyvinyl alcohol\*, povidone (K-30)\*, silicon dioxide, sodium starch glycolate, stearic acid, talc\*, titanium dioxide

\* may contain

## **IBUPROFEN**

***Questions or comments?1-800-634-7680***

## **DIAMODE**

### ***Active ingredient (in each caplet)***

Loperamide Hydrochloride 2 mg

## **DIAMODE**

### ***Purpose***

Antidiarrheal

## **DIAMODE**

### *Uses*

Controls the symptoms of diarrhea, including Traveler's diarrhea

## **DIAMODE**

### *Warnings*

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl

## **DIAMODE**

**Do not use** if you have bloody or black stool

## **DIAMODE**

**Ask a doctor before use if you have**

- a fever
- mucus in stool
- a history of liver disease

## **DIAMODE**

**Ask a doctor or pharmacist before use if you are** taking antibiotics

## **DIAMODE**

**When using this product**

- tiredness, drowsiness or dizziness may occur
- be careful when driving or operating machinery

## **DIAMODE**

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

## **DIAMODE**

**If you are pregnant or breast-feeding,** ask a health professional before use

## **DIAMODE**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

## **DIAMODE**

### *Directions*

- do not use more than directed

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

---

<b>Adults and children: (12 years and older)</b>	Take 2 caplets after the first loose stool followed by 1 caplet after each subsequent loose stool but no more than 4 caplets in 24 hours.
<b>Children under 12 years:</b>	Do not give to children under 12 years of age.

---

## **DIAMODE**

### ***Other information***

- store at room temperature 68° - 77°F (20° - 25°C)
- tamper-evident sealed packets
- do not use any opened or torn packet

## **DIAMODE**

***Inactive ingredients*** anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow #10, FD&C Blue #1, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

## **DIAMODE**

***Questions or comments? 1-800-634-7680***

## **EXTRA STRENGTH NON-ASPIRIN**

**EXTRA STRENGTH**

**NON-ASPIRIN**

2 Tablets

*Mfd. for **MEDIQUE PRODUCTS • Fort Myers, FL 33967***



## ANTACID

## ANTACID

2 Tablets

Mfd for MEDIQUE PRODUCTS, Fort Myers, FL 33967



## COLD RELIEF

## COLD RELIEF

2 Tablets

Mfd for *MEDIQUE PRODUCTS, Fort Myers, FL 33967*



## IBUPROFEN

IBUPROFEN

2 Tablets

Mfd. for: *MEDIQUE PRODUCTS • Fort Myers, FL 33967*




## DIAMODE

Medique

**Diamode**

**1 Caplet**

Mfd. for *MEDIQUE PRODUCTS, Fort Myers, FL 33967*



## **OUTER CARTON PDP**

MEDS ON THE GO...

INSTANT RELIEF ON THE GO!

5 Different OTC Medications

17 Individual Packets

HEADACHE, TOOTHACHE, MUSCLE ACHES, DIARRHEA, UPSET STOMACH, INDIGESTION, COUGH/COLD...AND MORE!

Acetaminophen • Ibuprofen • Anti-Diarrheal • Antacid • Decongestant



# MEDS ON THE GO...

**INSTANT RELIEF ON THE GO!**

**5** Different OTC Medications

**17** Individual Packets

**HEADACHE, TOOTHACHE, MUSCLE ACHES, DIARRHEA, UPSET STOMACH, INDIGESTION, COUGH/COLD... AND MORE!**

Acetaminophen • Ibuprofen • Anti-Diarrheal • Antacid • Decongestant

## MEDS ON THE GO

acetaminophen, calcium carbonate, dextromethorphan hbr, guaifenesin, phenylephrine hcl, ibuprofen, loperamide hcl kit

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:72082-002
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72082-002-05	1 in 1 BOX; Type 1: Convenience Kit of Co-Package	01/01/2019	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 PACKET	3 in 2
Part 2	2 PACKET	4 in 2
Part 3	2 PACKET	3 in 2
Part 4	2 PACKET	3 in 2
Part 5	4 PACKET	4

### Part 1 of 5

# MEDI-FIRST NON-ASPIRIN EXTRA STRENGTH

acetaminophen tablet, film coated

## Product Information

**Item Code (Source)** NDC:47682-126

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

## Product Characteristics

<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (Round)	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	FR;33
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-126-99	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/01/2019	

# MEDI-FIRST ANTACID

calcium carbonate tablet, chewable

## Product Information

Item Code (Source)	NDC:47682-820
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB, CARBONATE ION - UNII:7UJQ5OPE7D)	CALCIUM CARBONATE	420 mg

## Inactive Ingredients

Ingredient Name	Strength
MALTO DEXTRIN (UNII: 7CVR7L4A2D)	
SUCROSE (UNII: C151H8M554)	
ASPARTAME (UNII: Z0H242BBR1)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
ACACIA (UNII: 5C5403N26O)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MINERAL OIL (UNII: T5L8T28FGP)	
SORBITOL (UNII: 506T60A25R)	

## Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	AZ;036
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-820-99	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part331	01/01/2019	

# MEDI-FIRST COLD RELIEF

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride tablet

## Product Information

Item Code (Source) NDC:47682-139

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (round)	Size	12mm
Flavor		Imprint Code	FR;12
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-139-99	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	01/01/2019	

# MEDI-FIRST IBUPROFEN

ibuprofen tablet, coated

## Product Information

Item Code (Source) NDC:47682-708

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE K30 (UNII: U725QWY32X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	

## Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G;2
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-708-99	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/01/2019	

## Part 5 of 5

### MEDIQUE DIAMODE

loperamide hydrochloride tablet

## Product Information

Item Code (Source)	NDC:47682-200
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4I)	LOPERAMIDE HYDROCHLORIDE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CROSPVIDONE (UNII: 2S7830E561)	
CORN OIL (UNII: 8470G57WFM)	
POWDERED CELLULOSE (UNII: SMD1X3XO9M)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	

## Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	123
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-200-46	1 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	01/01/2019	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/01/2019	

**Labeler** - Doc in the Box LLC (081033259)

## Establishment

Name	Address	ID/FEI	Business Operations
Doc in the Box LLC		081033259	repack(72082-002)

Revised: 1/2019

Doc in the Box LLC