

**4376 FIRST AID KIT- 4376 first aid kit  
Honeywell Safety Products USA, INC**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**0498-4376: First Aid Kit (BZK wipe,FABC, neomycin, alcohol wipes, sting relief, ASA, aypanal- Z019807)**

**Alcohol Wipe**  
***Active ingredient***

Isopropyl alcohol 70%

**Alcohol Wipe**  
***Purpose***

First aid antiseptic

**Alcohol Wipe**  
***Uses***

- first aid to help prevent infection in minor cuts, scrapes, and burns

**Alcohol Wipe**  
***Warnings***

**For external use only**

**Do not use**

- in the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burn

**When using this product**

- do not use longer than one week unless directed by a doctor

**Stop use and consult a doctor**

- if condition persists or gets worse

**Keep out of reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away.

**Alcohol Wipe*****Directions***

- clean the affected area
- apply wipe to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard wipe after single use

**Alcohol Wipe*****Other information***

store at room temperature 15 ° to 25 ° C (59 ° to 77 °F)

**Alcohol Wipe*****Inactive ingredient***

water

**Alcohol Wipe*****Questions***

1-800-430-5490

**Aspirin*****Active ingredient (in each tablet)***

Aspirin 325 mg (NSAID)\* \*nonsteroidal anti-inflammatory drug

**Aspirin*****Purpose***

Pain reliever/fever reducer

**Aspirin*****Uses***

temporarily reduces fever and relieves minor aches and pains associated with:

- a cold
- headache
- toothache
- muscular aches
- backache
- minor pain of arthritis
- premenstrual and menstrual periods

## **Aspirin**

### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:are:

- age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout or arthritis

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- ringing in the ears or loss of hearing occurs
- any new symptoms appear

**If pregnant or breast-feeding,**

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

- In case of overdose, get medical help or contact Poison Control Center right away.

**Aspirin**

***Directions***

- drink a full glass of water with each dose
- adults and children 12 years of age and older: take 1 or 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- children under 12 years of age: consult a doctor

**Aspirin**

***Other information***

- store at room temperature 15° - 30°C (59° - 86°F)
- TAMPER EVIDENT PACKETS
- DO NOT USE IF OPEN OR TORN

**Aspirin**

***Inactive ingredients***

corn starch, croscarmellose sodium\*, hypromellose\*, microcrystalline cellulose\*, mineral oil\*, polyethylene glycol\*, povidone, propylene glycol, silicon dioxide, stearic acid\*, titanium dioxide\*

\*may contain these ingredients

**Aspirin**

***Questions or Comments***

1-800-430-5490

**Sting Relief**

***Active ingredient***

Ethyl alcohol 50.0%

Lidocaine HCl 2.0%

**Sting Relief**

## **Purpose**

Antiseptic

Topical pain relief

## **Sting Relief**

### **Uses**

- prevent infection in minor scrapes, and temporary relief of itching of insect bites

## **Sting Relief**

### **Warnings**

**For external use only**

**Flammable, keep away from open fire or flame**

### **Do not use**

- over large areas of the body
- in eyes
- over raw or blistered areas

### **Stop use and ask a doctor**

- if conditions worsen or persist for more than 7 days or clear up and occur again within a few days

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

## **sting Relief**

### **Directions**

- adults and children 2 years and older: Apply to cleaned affected area not more than 3 times daily.
- children under 2 years of age: consult a doctor.

## **Sting Relief**

### **Inactive ingredients**

benzalkonium chloride, menthol, and purified water

## **Sting Relief**

### **Questions or Comments?**

1-800-430-5490

## **BZK**

### **Active ingredient**

Benzalkonium chloride 0.13% w/v

## **BZK**

### ***Purpose***

First aid antiseptic

## **BZK**

### ***Uses***

Antiseptic cleansing of face, hands, and body without soap and water

## **BZK**

### ***Warnings***

#### **For external use only**

#### **Do not use**

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor
- longer than 1 week unless directed by a doctor

#### **Stop use and ask a doctor if**

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

#### **Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **BZK**

### ***Directions***

- tear open packet and use as a washcloth

## **BZK**

### ***Other information***

- store at room temperature 15 ° to 30 ° C (59 ° - 86 ° F)
- do not reuse towelette

## **BZK**

### ***Inactive ingredient***

water

**BZK**

**Questions**

1-800-430-5490

**Neomycin**

**Active ingredient (each gram contains)**

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

**Neomycin**

**Purpose**

First aid antibiotic

**Neomycin**

**Uses**

- first aid to help prevent infection in
- minor cuts
- scrapes
- burns

**Do not use**

- in the eyes
- over large areas of the body

**Neomycin**

**Warnings**

**For external use only**

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**Stop use and ask a doctor if**

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away

**Neomycin**

**Directions**

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

## **Neomycin**

### ***Other information***

- store at 15<sup>0</sup> to 25<sup>0</sup> C (59<sup>0</sup> to 77<sup>0</sup> F)

## **Neomycin**

### ***Inactive ingredient***

petrolatum

## **Neomycin**

### ***Questions***

1-800-430-5490

## **Aypanal EX**

### ***Active ingredient***

Acetaminophen 500 mg

## **Aypanal EX**

### ***PURpose***

Pain reliever/fever reducer

## **Aypanal EX**

### ***Uses***

- temporarily relieves minor aches and pains due to the common cold and headache - temporarily reduces fever

## **Aypanal EX**

### ***Warnings***

**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount.
- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If skin reaction occurs, stop use and seek medical help right away

### **Do Not Use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask a doctor before use if you have**

- liver disease

### **Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present
- If pregnant or breastfeeding,
- ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Aypanal EX**

#### ***Directions***

- **do not take more than directed (see overdose warning)**
- adults and children 12 years of age and over: Take 2 tablets with water every 6 hours while symptoms last.
- do not take any more than 8 tablets in 24 hours.
- children under 12: consult a doctor

### **Aypanal EX**

#### ***Other information***

- store at room temperature 15<sup>o</sup> -30<sup>o</sup> C (59<sup>o</sup> -86<sup>o</sup> F)
- TAMPER EVIDENT- DO NOT USE IF OPEN OR TORN

### **Aypanal EX**

#### ***Inactive ingredients***

microcrystalline cellulose, povidone, sodium starch glycolate, starch, stearic acid

## **Aypanal EX**

### ***Questions or Comments?***

1-800-430-5490

## **FABC**

### ***Active ingredient***

Benzalkonium chloride 0.13%

Lidocaine HCl 0.5%

## **FABC**

### ***Purpose***

First Aid antiseptic

External analgesic

## **FABC**

### ***Uses***

- prevent skin infection
- for temporary relief of pain associated with minor burns

## **FABC**

### ***Warnings***

For external use only

### **Do not use**

- in or near the eyes
- if you are allergic to any of the ingredients
- in large areas of the body, particularly over raw surfaces or blistered areas
- for more than 10 days

### **Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

### **Stop use and ask a doctor if**

- condition worsens
- symptoms persist for more than 7 days or clear up and occurs again within a few days

### **Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away.

## **FABC**

### ***Directions***

- **adults and children 2 years of age and older:**
- clean the affected area
- apply a small amount of this product (equal to the surface area of the tip of a finger) onto affected area 1 to 3 times daily
- may be covered with a sterile bandage
- **children under 2 years of age: consult a doctor**

## **FABC**

### ***Other information***

- tamper evident sealed packets
- do not use if packet is opened or torn

## **FABC**

### ***Inactive ingredients***

aloe barbadensis juice, cetyl alcohol, diazolidinyl urea, edetate disodium, glycerin, glyceryl stearate SE, methylparaben, mineral oil, PEG-100, propylene glycol, propylparaben, stearic acid, trolamine, water

## **FABC**

### ***Questions***

1-800-430-5490

## **4376**


### **Z019807 KIT CONTENTS**

1 ADHESIVE TAPE W/P 1/2"X 5 YD  
1 FIRST AID GUIDE ASHI  
1 EMERGENCY SURVIVAL BLANKET  
1 GAUZE CLEAN-WRAP BDGE N/S 2"  
10 CTA 6" SINGLE TIP 100/BAG  
1 SCISSOR BDGE 4" RED PLS HDL  
1 FANNY PACK RED FAK LOGO EMPTY  
4 TONGUE BLADES SR WRAPPED 250  
LBL STOCK 6-3/8"X4"  
1 LBL STOCK 3"x1-7/8"

1 LBL CONTS 6 3/4"X3 1/2" ID B  
6 BZK ANTISEPTIC WIPE, BULK  
1 PR LRG NITRILE GLVES ZIP BAG  
3 FIRST AID BURN CREAM 1.0GR PKT EACH  
3 POUCH NEOMYCIN ANTIBIOTIC .9 G  
4 ADH BANDAGE BUTTERFLY 1980000  
6 WIPE ALCOHOL PREP IPA 70% (DUKAL)  
2 LANCET BLADE #11 STERILE  
10 ADH BANDAGE 3/8" X 1 1/2" DNX  
3 SAFETEC STING RELIEF WIPES BULK  
1 COLD PACK UNIT 4"X6" BULK  
4 GAUZE PADS 2"X2" 12PLY  
2 GAUZE PADS 4"X4" 12PLY  
15 PLASTIC BANDAGE 3/4" X 3"  
1 WOVEN KNUCKLE BANDAGE  
1 ZIP-LOCK BAG 5" X 5" .002  
2 AYPANAL BULK 2/PK  
2 ASPIRIN BULK 2/PK

### **Alcohol Wipe Label**

825362T Rev. A  
 white printed  
 Pms 072 blue, Pms 186 red,  
 Pms 7459 blue and black



## Aspirin Principal Display Panel



# Sting Relief Principal Display Panel

825366 Rev B

Honeywell

**032043P**




**Sting Relief Wipes**

**Use for:**  
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches  
Saturated Wipes

100 wipes

**032043P**



**Sting Relief Wipes**

**Use for:**  
Minor Cuts • Scrapes • Insect Bites

Single Use Pouches  
Saturated Wipes

100 wipes

**Drug Facts**

Active Ingredients	Purpose
Ethyl alcohol 50.0%	First aid antiseptic
Lidocaine HCl 2.0%	Topical analgesic

**Uses** First aid to help prevent infection in minor scrapes and temporary relief of itching of insect bites.

**Warnings**  
For external use only.  
Flammable, keep away from fire or flame.

**Do not use** • over large areas of the body • in eyes • over raw or blistered areas

**Stop use and ask a doctor** • if conditions worsen or persist for more than 7 days or clear up and occur again within a few days.

**Keep out of reach of children.**  
If swallowed get medical help or contact Poison Control center right away.

**Directions**  
• adults and children 2 years and older, apply to cleaned affected area not more than 3 times daily.  
• children under 2 years of age, consult a doctor.

**Inactive Ingredients**  
benzalkonium chloride, menthol, purified water

**Questions or comments?** 1-800-430-5490

Distributed by  
Honeywell Safety  
Products, USA, Inc.  
Smithfield, RI 02917

[www.honeywellsafety.com](http://www.honeywellsafety.com)

USA  
1-800-430-5490  
0-401-943-4400



8 21812 01284 1

Made in USA  
Packaged in Mexico

# BZK Principal Display Panel

47001083  
Rev B

Antiseptic Towelettes

**Honeywell**

02-16-35MD

**Antiseptic Towelettes**

*Benzalkonium chloride*  
*First aid antiseptic*

**Six-Saturated Towelettes**

Distributed by  
Honeywell Safety  
Products USA, Inc.  
Smithfield, RI 02917

47001083  
Rev B

**Antiseptic Towelettes**

Honeywell

**Drug Facts**

Active Ingredient	Purpose
Benzalkonium chloride 0.133% w/v	First aid antiseptic

**Uses** • antiseptic cleaning of face, hands and body without soap and water. • air dries in seconds

**Warnings**  
For external use only

**When using this product** • do not use in the eyes or apply over large areas of the body  
**Ask a doctor before use** • in case of deep or puncture wounds, animal bites, or serious burns

**Stop use and consult a doctor if**  
• irritation, redness or other symptoms develop • condition persists or gets worse

**Do not use** • longer than 1 week unless directed by doctor

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** • tear open packet, unfold and use as washcloth



**Other information**  
• store at room temperature 15° -30° C (59° -86° F) • do not reuse towelette

**Inactive ingredient** water

**Questions or comments** 1-800-430-5490

**Neomycin**  
**Principal Display Panel**

796041-25 Rev A Unit Carton Printing Plate for "C" size carton.





901850-10

**First Aid Burn Cream**

021020-10

**First Aid Burn Cream**

*Benzalkonium chloride*

*First aid antiseptic*

*Lidocaine HCl*

*External analgesic*

Distributed by  
Honeywell Safety  
Products USA, Inc.  
Smithfield, RI 02917

10 Packets, Net Wt 1/32 oz (0.9g) each

**First Aid Burn Cream**

901850-10

<b>Drug Facts</b>	
<b>Active ingredients</b> Benzalkonium chloride 0.13% Lidocaine HCl 0.5%	<b>Purposes</b> First aid antiseptic External analgesic
<b>Use</b> - prevent skin infection - for the temporary relief of pain associated with minor burns	
<b>Warnings For external use only</b>	
<b>Do not use</b> - in or near the eyes - if you are allergic to any of the ingredients - in large areas of the body particularly over raw surfaces or blistered areas - for more than 10 days	
<b>Ask a doctor before use if you have</b> - deep or puncture wounds - animal bites - serious burns	
<b>Stop use and ask a doctor if</b> - condition worsens - symptoms persist more than 7 days or clear up and occur again within a few days	
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away	
<b>Directions</b>	
<b>adults and children 2 years and over:</b> - clean the affected area - apply a small amount of this product (equal to the surface area of the tip of a finger) onto affected area 1 to 2 times daily - may be covered with a sterile bandage	
<b>children under 2 years:</b> ask a doctor	
<b>Other information</b> - tamper evident sealed packets - do not use if packet is opened or torn	
<b>Inactive ingredients</b> aloe barbadensis leaf juice, cetyl alcohol, diazolidinyl urea, edetate disodium, glycerin, glyceryl stearate SE, methylparaben, mineral oil, PEG-100, propylene glycol, propylparaben, stearic acid, triethylamine, water	
<b>Questions or comments?</b> 1-800-430-5430	

**4376 Kit Label  
Z019807**



Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917

**APPROVED**  
By Rodrigo Rosas Abiano at 2:58 pm, Mar 11, 2019

**4376 FIRST AID KIT**

4376 first aid kit kit

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0498-4376
---------------------	----------------	---------------------------	---------------

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4376-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

**Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
<b>Part 1</b>	3 PACKET	2.7 g
<b>Part 2</b>	6 POUCH	2.4 mL
<b>Part 3</b>	2 PACKET	4
<b>Part 4</b>	3 PACKET	2.7 g
<b>Part 5</b>	6 PACKET	8.4 mL

<b>Part 6</b>	3 POUCH	1.2 mL
<b>Part 7</b>	2 PACKET	4

## Part 1 of 7

### FIRST AID BURN

benzalkonium chloride, lidocaine hydrochloride cream

#### Product Information

<b>Item Code (Source)</b>	NDC:0498-0903
<b>Route of Administration</b>	TOPICAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	0.13 g in 100 g
<b>LIDOCAINE HYDROCHLORIDE</b> (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE	0.5 g in 100 g

#### Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>ALOE VERA LEAF</b> (UNII: ZY81Z83H0X)	
<b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)	
<b>PEG-100 STEARATE</b> (UNII: YD01N1999R)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>TROLAMINE</b> (UNII: 9O3K93S3TK)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XXE4)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>DIAZOLIDINYL UREA</b> (UNII: H5RIZ3MPW4)	

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		0.9 g in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/20/2017	

## Part 2 of 7

### ALCOHOL WIPE

isopropyl alcohol swab

## Product Information

Item Code (Source)	NDC:0498-0143
Route of Administration	TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

## Part 3 of 7

### ASPIRIN

aspirin tablet

## Product Information

**Item Code (Source)** NDC:0498-0114

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	FR21
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0114-01	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

## Part 4 of 7

### NEOMYCIN

antibiotic ointment

#### Product Information

Item Code (Source)	NDC:0498-0730
Route of Administration	TOPICAL

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g

#### Inactive Ingredients

Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0730-01	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

#### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/31/2010	

## Part 5 of 7

### ANTISEPTIC TOWELETTE

benzalkonium chloride liquid

#### Product Information

Item Code (Source)	NDC:0498-0501
Route of Administration	TOPICAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZALKONIUM CHLORIDE	1.3 mg in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0501-00	1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/22/2017	

**Part 6 of 7****STING RELIEF PAD**

ethyl alcohol, lidocaine swab

**Product Information**

<b>Item Code (Source)</b>	NDC:0498-0733
<b>Route of Administration</b>	TOPICAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>LIDOCAINE HYDROCHLORIDE</b> (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	20 mg in 1 mL
<b>ALCOHOL</b> (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M)	ALCOHOL	0.5 mL in 1 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>WATER</b> (UNII: 059QF0KO0R)	

MENTHOL (UNII: L7T10EIP3A)

BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0733-00	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/23/2017	

### Part 7 of 7

### AYPANAL EX

acetaminophen tablet

### Product Information

Item Code (Source)	NDC:0498-2110
Route of Administration	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

### Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	FR1

**Contains****Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-2110-01	2 in 1 PACKET; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/02/2017	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

**Labeler** - Honeywell Safety Products USA, INC (118768815)

Revised: 1/2024

Honeywell Safety Products USA, INC