

IBUPROFEN 200 MG- ibuprofen capsule, liquid filled
Allegiant Health

363,368 - Ibuprofen 200mg

Active ingredient(s)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)* (present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Use(s)

temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chances are higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma , or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if

- under a doctor’s care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

- any new symptoms appear

Pregnancy/Breastfeeding

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely

directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- **adults and children 12 years and over:** take 1 softgel every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 softgel, 2 softgels may be used
- do not exceed 6 softgels in 24 hours, unless directed by a doctor
- **children under 12 years:** ask a doctor

Other information

- each softgel contains: potassium 20mg
- store between 20-25 °C (68-77° F)
- read all warnings and directions before use
- do not use if seal under bottle cap is broken or missing.

Inactive ingredients

ammonium hydroxide, FD&C green #3 aluminum lake, gelatin, iron oxide black, medium chain triglycerides, polyethylene glycol, propylene glycol, potassium hydroxide, purified water, shellac, sorbitol sorbitan, sorbitan monooleate

Questions

Call 1-888-952-0050

Monday through Friday 9AM – 5PM EST.

Principal Display Panel

FERROSFERRIC OXIDE (UNII: XM0M87F357)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SHELLAC (UNII: 46N107B71O)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
SORBITOL SOLUTION (UNII: 8KW3E207O2)	

Product Characteristics

Color	green	Score	no score
Shape	CAPSULE	Size	15mm
Flavor		Imprint Code	133
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-368-07	1 in 1 CARTON	12/22/2014	
1		10 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69168-368-30	1 in 1 CARTON	12/22/2014	
2		30 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079205	12/22/2014	

Labeler - Allegiant Health (079501930)

Revised: 12/2021

Allegiant Health