

**ANTI DIARRHEAL- loperamide hcl tablet, film coated
H E B**

HEB Anti-Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

| | |
|--|--|
| adults and children 12 years and over | 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours |
| children 9-11 years (60-95 lbs) | 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours |
| children 6-8 years (48-59 lbs) | 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours |
| children 2-5 years (34 to 47 lbs) | ask a doctor |
| children under 2 years (up to 33 lbs) | do not use |

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Imodium® A-D active ingredient

See New Warnings

Anti-Diarrheal

Loperamide Hydrochloride Tablets, 2 mg

Anti-Diarrheal

Controls The Symptoms of Diarrhea

actual size

12 CAPLETS*

*Capsule-Shaped Tablets

Drug Facts (continued)
Inactive ingredients anhydrous lactose, carnauba wax, D & C yellow no. 10 aluminum lake, FD & C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Drug Facts (continued)
Questions or comments?
 1-800-719-9260
GLUTEN FREE

Drug Facts

Active ingredient
 (in each caplet)
 Loperamide HCl 2 mg.....Anti-diarrheal

Purpose
 Use controls symptoms of diarrhea, including Traveler's Diarrhea.

Warnings
 Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
 Heart alert: Taking more than directed can cause serious heart problems or death
 Do not use if you have bloody or black stool
 Ask a doctor before use if you have
 ■ fever ■ mucus in the stool
 ■ a history of liver disease
 ■ a history of abnormal heart rhythm
 Ask a doctor or pharmacist before use if you are taking a prescription drug.
 Loperamide may interact with certain prescription drugs.
 When using this product, dizziness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.
 Stop use and ask a doctor if
 ■ symptoms get worse
 ■ diarrhea lasts for more than 2 days
 ■ you get abdominal swelling or bloating
 These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

Other information
 ■ store at 20°-25°C (68°-77°F)
 ■ see end panel for lot number and expiration date

| | |
|---------------------------------------|--|
| adults and children 12 years and over | 2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours |
| children 6-8 years (48-59 lbs) | 1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours |
| children 2-5 years (34 to 47 lbs) | ask a doctor |
| children under 2 years (up to 33 lbs) | do not use |

Directions
 ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
 ■ find right dose on chart. If possible, use weight to dose: o the wise, use age.
 Keep out of reach of children. In case of overdose, get medical help or contact Poison Control Center right away (1-800-222-1222).

Drug Facts (continued)



13463-2001

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D.

OPEN OTHER END

GOODIE / A / R / H / A

22453 1J C8

MADE WITH PRIDE & CARE FOR H-E-B® SAN ANTONIO, TX 78204

H-E-B
 100% GUARANTEE promise
 If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

Compare to Imodium® A-D active ingredient**

NDC 37808-552-53

See New Warnings



Anti-Diarrheal



ANTI DIARRHEAL

loperamide hcl tablet, film coated

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:37808-552 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|--------------------------|----------|
| LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II) | LOPERAMIDE HYDROCHLORIDE | 2 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | |
| D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) | |
| FD&C BLUE NO. 1 (UNII: HBR47K3TBD) | |

Product Characteristics

| | | | |
|----------|-------|--------------|----------|
| Color | GREEN | Score | 2 pieces |
| Shape | OVAL | Size | 10mm |
| Flavor | | Imprint Code | L2 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:37808-552-23 | 1 in 1 CARTON | 05/01/2014 | |
| 1 | | 72 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 2 | NDC:37808-552-16 | 2 in 1 CARTON | 05/01/2014 | 03/31/2020 |
| 2 | | 72 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 3 | NDC:37808-552-67 | 1 in 1 CARTON | 06/18/2018 | |
| 3 | | 48 in 1 BOTTLE; Type 0: Not a Combination Product | | |
| 4 | NDC:37808-552-53 | 12 in 1 CARTON | 02/20/2020 | |
| 4 | | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA075232 | 05/01/2014 | |

Labeler - HEB (007924756)

Revised: 2/2020

HEB