

**ANTI-DIARRHEAL- loperamide hcl tablet  
H E B**

-----  
**HEB 44-375**

***Active ingredient (in each caplet)***

Loperamide HCl 2 mg

***Purpose***

Anti-diarrheal

***Use***

controls symptoms of diarrhea, including Travelers' Diarrhea

***Warnings***

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use**

if you have bloody or black stool.

**Ask a doctor before use if you have**

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product,**

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### **Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

### **Inactive ingredients**

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

### **Questions or comments?**

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

### **Principal Display Panel**

**Compare to Imodium® A-D** active ingredient\*

NDC 37808-375-08

**H-E-B®**

**Anti-Diarrheal**

Loperamide Hydrochloride 2 mg  
Anti-Diarrheal

**Diarrhea Relief**

Controls The Symptoms of Diarrhea,  
Including Travelers' Diarrhea

**24 CAPLETS**

Each caplet contains 2 mg loperamide hydrochloride

actual  
size

**1.** To open, tear  
at perforations and  
remove a section.

**2.** Use ✂ to cut  
through plastic  
for caplet.

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR  
SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by Johnson & Johnson Corporation,  
owner of the registered trademark Imodium® A-D.

50844 REV0619A37508

**MADE WITH PRIDE AND CARE FOR  
H-E-B®, SAN ANTONIO, TX 78204**

**H-E-B®**

100%  
GUARANTEE

*promise*

If you aren't completely pleased  
with this product, we'll be happy to  
replace it or refund your money.  
You have our word on it.

\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark "Imodium" A-D.

**Drug Facts (continued)**  
 ■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number  
**Inactive ingredients** corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dryhydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide  
**Questions or comments?** Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

**Other information**  
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool but no more than 2 caplets in 24 hours
children 2-5 years (24-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Directions** ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea  
 ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

**Warnings**  
 Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.  
 Heart alert: Taking more than directed can cause serious heart problems or death.  
 Do not use if you have bloody or black stool.  
 Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm  
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.  
 When using this product, dizziness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.  
 Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bloating. These may be signs of a serious condition.  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Use** controls symptoms of diarrhea, including travelers' diarrhea  
**Active ingredient (in each caplet)** Loperamide HCl 2 mg  
**Purpose** Anti-diarrheal

**Drug Facts** KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION



**MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204**

**100% GUARANTEE**  
 If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

No print/No varnish Lot & Exp date

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING**



# Anti-Diarrheal

Loperamide Hydrochloride 2 mg/Anti-Diarrheal

Diarrhea Relief

Compare to *Imodium*® A-D active ingredient\*


NDC 37808-375-08



# Anti-Diarrheal

Loperamide Hydrochloride 2 mg

B-0712-375-08-R  
 REV0619A37508



44-375

## ANTI-DIARRHEAL

loperamide hcl tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37808-375
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;375
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-375-08	4 in 1 CARTON	05/03/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:37808-375-02	2 in 1 CARTON	05/03/2005	02/18/2022
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/03/2005	

**Labeler** - H E B (007924756)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(37808-375) , pack(37808-375)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(37808-375)

Revised: 3/2024

H E B