

DIGESTIVE RELIEF REGULAR STRENGTH- bismuth subsalicylate liquid
TARGET Corporation

Drug Facts

Active ingredient (in each 30 mL)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before using
- mL = milliliter
- measure only with dosing cup provided. do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL (1 dose) every 1/2 to 1 hour as needed
 - do not exceed 8 doses (240 mL) in 24 hours
 - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** potassium 8 mg sodium 9 mg
- **each 30 mL contains:** salicylate 243 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid

Questions?

Call **1-800-910-6874**

Principal Display Panel

Compare to the active ingredient in Pepto-Bismol®*

5-symptom

digestive relief

Bismuth subsalicylate 525 mg per 30 mL

upset stomach reliever/antidiarrheal

Soothing relief of heartburn, indigestion, nausea,

upset stomach and diarrhea

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Dist. by Target Corp.,

Mpls., MN 55403

Package Label

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ mL = milliliter ■ shake well before using ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 30 mL (1 dose) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years of age: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each 30 mL contains: potassium 8 mg, sodium 9 mg ■ each 30 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments? Call 1-800-910-6874

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

245 05 0589 R00
C-001744-01-045
Dist. by Target Corp.
Mpls., MN 55403
TM & ©2022
Target Brands, Inc.



Compare to active ingredient in Pepto-Bismol® Regular Strength*

5-symptom digestive relief

bismuth subsalicylate 525 mg upset stomach reliever / antidiarrheal

relieves nausea, heartburn, indigestion, upset stomach and diarrhea



NDC 11673-372-08

Drug Facts TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Active ingredient (in each 30 mL dose) Purpose Bismuth subsalicylate 525 mg.....Upset stomach reliever/Antidiarrheal

Uses relieves ■ diarrhea ■ travelers' diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ gas ■ indigestion ■ nausea ■ fullness ■ belching

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are: ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

TARGET Digestive Relief

DIGESTIVE RELIEF REGULAR STRENGTH

bismuth subsalicylate liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11673-372 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-----------------------|-----------------|
| BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ) | BISMUTH SUBSALICYLATE | 525 mg in 30 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| BENZOIC ACID (UNII: 8SKN0B0MIM) | |
| D&C RED NO. 22 (UNII: 1678RKX8RT) | |
| D&C RED NO. 28 (UNII: 767IP0Y5NH) | |
| MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC) | |
| WATER (UNII: 059QF0KOOR) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| SALICYLIC ACID (UNII: O414PZ4LPZ) | |
| SODIUM SALICYLATE (UNII: WQ1H85SYP) | |
| SORBIC ACID (UNII: X045WJ989B) | |
| METHYLCELLULOSE (1500 CPS) (UNII: PONTE48364) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:11673-372-16 | 473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 07/31/2018 | |
| 2 | NDC:11673-372-08 | 236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 07/31/2018 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M008 | 07/31/2018 | |

Labeler - TARGET Corporation (006961700)

Revised: 6/2024

TARGET Corporation