

PEPTO RELIEF- bismuth subsalicylate tablet, chewable
Advance Pharmaceutical Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

PEPTO RELIEF Bismuth Subs alicylate 262 MG

Drug Facts

.

Active Ingredient

(in each tablet)

Bismuth subsalicylate 262 mg

Purpose

Anti-diarrheal/Upset stomach reliever

Uses

- Controls diarrhea
- Relieves upset stomach due to overindulgence in food or drink

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

- If you have bloody or black stool
- If you have an ulcer or bleeding problem

Ask a doctor before use if you have

- Fever
- Mucus in the stool

Ask a doctor or pharmacist before use if you are

- taking any drug for
- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

when using this product a temporary, but harmless, darkening of the stool and/or tongue may occur
stop use and ask a doctor if

- symptoms get worse

- ringing in the ears or loss of hearing occurs

diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- repeat dosage every 1/2 to 1 hour as needed
- do not take more than 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- **adults and children 12 years and over:** 2 tablets
- **children under 12 years:** ask a doctor

Other Information

- **Each tablet contains:** calcium 77 mg, salicylate 102 mg
- store at 15°-30°C (59°-86°F)
- protect from moisture

INACTIVE INGREDIENT

calcium carbonate, D&C red # 27 (Al-lake), dextrose, flavor (cherry), magnesium stearate, maltodextrin, silicon dioxide, sorbitol

Questions or Comments

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Call 631-981-4600, 8.30 am – 4.30 pm EST Monday - Friday

Package Label



Advance
Pharmaceutical Inc.

30 CHEWABLE TABLETS

Protective Coating Action
Soothing Relief for Upset Stomach, Diarrhea,
Indigestion, Heartburn and Nausea

UPSET STOMACH RELIEVER

ANTI-DIARRHEAL/

Bismuth Subsalicylate 262 mg

Pepto Relief

Compare to Active Ingredient in
PEPTO-BISMOL®*

NDC 17714-045-30

Drug Facts

Active ingredient (in each tablet)

Bismuth subsalicylate
262 mg

Purposes

Anti-diarrheal/
Upset stomach reliever

Uses

- controls diarrhea
- relieves upset stomach due to overindulgence in food or drink

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

- if you have bloody or black stool
- if you have an ulcer or bleeding problem

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*Advance Pharmaceutical, Inc. is not affiliated with the owner of the trademark PEPTO-BISMOL®

Manufactured by: Advance Pharmaceutical, Inc.
Ronkonkoma, NY 11779, U.S.A.



0 17714-045-30 2

LA1209



Lot No.:
Exp. Date:

Drug Facts (continued)

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- repeat dosage every 1/2 to 1 hour as needed
- do not take more than 8 doses in 24 hours
- use until diarrhea stops but not more than 2 days
- adults and children 12 years and over: 2 tablets
- children under 12 years: ask a doctor

Other information

- each tablet contains: calcium 77 mg, salicylate 102 mg
- store at 15°-30°C (59°-86°F)
- protect from moisture

Inactive ingredients calcium carbonate, D&C red #27 (Al-lake), dextrose, flavor (cherry), magnesium stearate, maltodextrin, silicon dioxide, sorbitol

Questions or comments?

call 631-981-4600, 8:30 am - 4:30 pm ET, Monday-Friday

NDC: 17714-045-30 – 30 CHEWABLE TABLETS

PEPTO RELIEF

bismuth subsalicylate tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:17714-045
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
DEXTROSE (UNII: IY9XDZ35W2)	
CHERRY (UNII: BUC5I9595W)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	pink (light)	Score	no score
Shape	ROUND	Size	16mm
Flavor	CHERRY	Imprint Code	AP;045
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17714-045-30	30 in 1 BOTTLE; Type 0: Not a Combination Product	01/09/2002	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	01/09/2002	

Labeler - Advance Pharmaceutical Inc. (078301063)

Registrant - Advance Pharmaceutical Inc. (078301063)

Establishment

Name	Address	ID/FEI	Business Operations
Advance Pharmaceutical Inc.		078301063	manufacture(17714-045)

Revised: 12/2017

Advance Pharmaceutical Inc.