

STOMACH RELIEF ULTRA- bismuth subsalicylate liquid
AmerisourceBergen (Good Neighbor Pharmacy) 46122

Drug Facts

Active ingredient (in each 15 mL)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adults and children 12 years and over:
 - 15 mL (1 dose) every 1/2 or 30 mL (2 doses) every hour as needed
diarrhea/traveler's diarrhea
 - 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

Other information

- **each 15 mL contains:** sodium 5 mg
- **each 15 mL contains:** salicylate 206 mg
- low sodium
- keep tightly closed

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Ultra Pepto-Bismol® active ingredient*

Ultra

Stomach Relief

2x Strength Per Ounce†

Upset Stomach Reliever/Antidiarrheal

(Bismuth Subsalicylate 525 mg)

Soothing Relief For 5 Symptoms:

- Upset Stomach
- Indigestion
- Heartburn
- Diarrhea
- Nausea

Alcohol Free

Sugar Free

FL OZ (mL)

†vs Original Pepto-Bismol®

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Distributed By AmerisourceBergen

1 West First Avenue, Conshohocken, PA 19428

www.mygnp.com

Package Label

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.
 In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ do not take more than 8 doses (120 mL) in 24 hours
 ■ use until diarrhea stops but not more than 2 days ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ mL = milliliter
 ■ shake well before using ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 15 mL (1 dose) every ½ hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 15 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ children under 12 years of age: ask a doctor

Other information ■ each 15 mL contains: sodium 5 mg
 ■ each 15 mL contains: salicylate 206 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, glycerin, purified water, sucralose, xanthan gum

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

†vs. Original Pepto-Bismol®

Distributed By AmerisourceBergen
 1 West First Avenue, Conshohocken, PA 19428
 Questions or Concerns? Visit us at www.mygnp.com



Compare to Ultra Pepto-Bismol® active ingredient*

NDC 46122-639-34

Ultra Stomach Relief

2x Strength Per Ounce!

Upset Stomach Reliever/Antidiarrheal
 (Bismuth Subsalicylate 525 mg)

Soothing Relief For 5 Symptoms:

- Upset Stomach • Indigestion
- Heartburn • Diarrhea • Nausea

Alcohol Free
 Sugar Free

8 fl oz (237 mL)

Drug Facts

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

Active ingredient (in each 15 mL dose)

Bismuth subsalicylate 525 mg.....Upset stomach reliver/Antidiarrheal

Purpose

Uses

- relieves
- travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn
 - indigestion ■ nausea ■ gas ■ fullness ■ belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are
 ■ allergic to salicylates (including aspirin)
 ■ taking other salicylate products

Do not use if you have

- an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have

- fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are

- taking any drug for ■ diabetes ■ gout ■ arthritis
- anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

GOOD NEIGHBOR PHARMACY Ultra Stomach Relief

STOMACH RELIEF ULTRA

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-639
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 7671P0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-639-34	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/06/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	07/06/2020	

Labeler - AmerisourceBergen (Good Neighbor Pharmacy) 46122 (007914906)

Revised: 2/2024

AmerisourceBergen (Good Neighbor Pharmacy) 46122