

**ALKA-SELTZER PLUS COUGH, MUCUS AND CONGESTION DAY AND NIGHT  
POWERMAX- acetaminophen, dextromethorphan hydrobromide, doxylamine  
succinate, guaifenesin, phenylephrine hydrochloride  
Bayer HealthCare LLC.**

-----  
**Alka-Seltzer Plus Maximum Strength Cough, Mucus & Congestion Day and  
Night Liquegels UI 1614294 and 1613941 24 count**

***Drug Facts***

**Product name**

**Do not take these products at the same time.**

**Alka-Seltzer Plus® Maximum Strength Cough, Mucus & Congestion Day  
PowerMax® Gels**

**Active ingredients and purpose**

***Active ingredients (in each capsule) Purposes***

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan hydrobromide 10 mg.....Cough suppressant  
Guaifenesin 200 mg.....Expectorant  
Phenylephrine hydrochloride 5 mg.....Nasal decongestant

**Purpose**

**Uses**

***Uses***

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves these symptoms due to a cold or flu:
  - nasal congestion · sinus congestion and pressure
  - minor aches and pains · headache
  - cough · sore throat
  - temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur

if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening · blisters · rash · hives
- facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

### **Ask a doctor before use if you have**

### **Ask a doctor before use if you have**

- liver disease ● heart disease ● high blood pressure
- thyroid disease ● diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

- cough with excessive phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**When using this product**

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

***Directions***

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after the last Night dose before starting Day product
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 6 capsules in 12 hours or as directed by a doctor.
- children under 12 years: do not use

**Other information**

## **Other information**

· store at 15°- 25°C (59° - 77°F)

## **Inactive ingredients**

**Inactive ingredients** FD&C red No. 40, gelatin, glycerin, lecithin, medium chain triglycerides, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

## **Questions or comments**

**Questions or comments? 1-800-986-0369** (Mon-Fri 9AM - 5PM EST)

## **Product name**

**Alka-Seltzer Plus® Maximum Strength Cough, Mucus & Congestion Night PowerMax® Gels**

## **Active ingredients and purpose**

### **Active ingredients (in each capsule) Purposes**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan hydrobromide 10 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine  
Phenylephrine hydrochloride 5 mg.....Nasal decongestant

## **Uses**

### **Uses**

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains · headache
- nasal and sinus congestion · cough
- sore throat · runny nose · sneezing
- temporarily reduces fever

## **Warnings**

### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening · blisters · rash · hives
- facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use to sedate children.**

**Do not use**

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

**Ask a doctor before use if you have**

**Ask a doctor before use if you have**

- liver disease ● heart disease ● high blood pressure
- thyroid disease ● diabetes ● glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are**

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

**When using this product**

- **do not exceed recommended dosage**
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children**

**Keep out of reach of children.** In case of overdose, get medical help

or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

## ***Directions***

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after the last Day dose before starting Night product
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 4 capsules in 12 hours or as directed by a doctor.
- children under 12 years: do not use

## **Other information**

### ***Other information***

- store at 15° - 25°C (59° - 77°F)

## **Inactive ingredients**

***Inactive ingredients*** D&C yellow No. 10, FD&C blue No. 1, gelatin, glycerin, polyethylene glycol, potassium aluminum silicate, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

## **Questions or comments**

***Questions or comments? 1-800-986-0369*** (Mon-Fri 9AM - 5PM EST)

## **Product carton label 24 count**

### **NEW**

Alka-Seltzer

PLUS

MAXIMUM

STRENGTH

Cough,

Mucus &

Congestion

POWERMAX® GELS

CONCENTRATED FORMULA

**DAY NON DROWSY**

**ACETAMINOPHEN / Pain Reliever-Fever Reducer**

Dextromethorphan HBr / Cough Suppressant

Guaifenesin / Expectorant

Phenylephrine HCl / Nasal Decongestant

- Nasal/Chest Congestion
- Headache, Body Ache Sore Throat, Fever
- Cough
- Mucus

16 LIQUID GELS

**NIGHT**

**ACETAMINOPHEN / Pain Reliever-Fever Reducer**

Dextromethorphan HBr / Cough Suppressant

Doxylamine Succinate / Antihistamine

Phenylephrine HCl / Nasal Decongestant

- Nasal / Chest Congestion
- Headache, Body Ache, Sore Throat, Fever
- Runny Nose, Sneezing
- Cough

8 LIQUID GELS: 24 TOTAL



## ALKA-SELTZER PLUS COUGH, MUCUS AND CONGESTION DAY AND NIGHT POWERMAX

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, guaifenesin, phenylephrine hydrochloride kit

### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:0280-0035

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0035-01	1 in 1 CARTON; Type 0: Not a Combination Product	06/01/2020	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 BLISTER PACK	6
Part 2	1 BLISTER PACK	2

## Part 1 of 2

### ALKA-SELTZER PLUS MAXIMUM STRENGTH COUGH, MUCUS AND CONGESTION DAY POWERMAX

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride capsule, liquid filled

## Product Information

Route of Administration ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
POTASSIUM ALUMINUM DISILICATE (UNII: SRB14JRX6C)	

<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	no score
<b>Shape</b>	OVAL (elliptical)	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	ASP;S
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		3 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2020	

## Part 2 of 2

### ALKA-SELTZER PLUS MAXIMUM STRENGTH COUGH, MUCUS AND CONGESTION NIGHT POWERMAX

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride capsule, liquid filled

### Product Information

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>GLYCERIN</b> (UNII: PDC6A3C00X)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>POTASSIUM ALUMINUM DISILICATE</b> (UNII: SRB14JRX6C)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL (elliptical)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	ASP;N
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		3 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2020	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/01/2020	

**Labeler** - Bayer HealthCare LLC. (112117283)