

BENZTROPINE MESYLATE- benztropine mesylate tablet
Chartwell RX, LLC.

Benztropine Mesylate Tablets

BENZTROPINE MESYLATE


TABLETS, USP

Rx Only

DESCRIPTION

Benztropine mesylate is a synthetic compound containing structural features found in atropine and diphenhydramine.

It is designated chemically as 3 α -(Diphenylmethoxy)-1 α H, 5 α H-tropane methanesulfonate. Its molecular formula is C₂₁ H₂₅ NO • CH₄ O₃ S, and its structural formula is:



Benztropine mesylate is a crystalline white powder, very soluble in water, and has a molecular weight of 403.54.

Each benztropine mesylate tablet for oral administration contains benztropine mesylate, USP, 0.5 mg, 1 mg, or 2 mg.

Each tablet contains the following inactive ingredients: dibasic calcium phosphate (hydrous), anhydrous lactose, magnesium stearate, microcrystalline cellulose, and pregelatinized starch.

CLINICAL PHARMACOLOGY

Benztropine mesylate possesses both anticholinergic and antihistaminic effects, although only the former have been established as therapeutically significant in the management of parkinsonism.

In the isolated guinea pig ileum, the anticholinergic activity of this drug is about equal to that of atropine; however, when administered orally to unanesthetized cats, it is only about half as active as atropine.

In laboratory animals, its antihistaminic activity and duration of action approach those of pyrilamine maleate.

INDICATIONS AND USAGE

Benztropine Mesylate Tablets, USP are indicated for use as an adjunct in the therapy of all forms of parkinsonism.

Useful also in the control of extrapyramidal disorders (except tardive dyskinesia-see **PRECAUTIONS**) due to neuroleptic drugs (e.g., phenothiazines).

CONTRAINDICATIONS

Hypersensitivity to benztropine mesylate tablets or to any component of the tablets.

Because of its atropine-like side effects, this drug is contraindicated in pediatric patients under three years of age, and should be used with caution in older pediatric patients.

WARNINGS

Safe use in pregnancy has not been established.

Benztropine mesylate may impair mental and/or physical abilities required for performance of hazardous tasks, such as operating machinery or driving a motor vehicle.

When benztropine mesylate is given concomitantly with phenothiazines, haloperidol, or other drugs with anticholinergic or antidopaminergic activity, patients should be advised to report gastrointestinal complaints, fever or heat intolerance promptly. Paralytic ileus, hyperthermia and heat stroke, all of which have sometimes been fatal, have occurred in patients taking anticholinergic-type antiparkinsonism drugs, including benztropine mesylate, in combination with phenothiazines and/or tricyclic antidepressants.

Since benztropine mesylate contains structural features of atropine, it may produce anhidrosis. For this reason, it should be administered with caution during hot weather, especially when given concomitantly with other atropine-like drugs to the chronically ill, the alcoholic, those who have central nervous system disease, and those who do manual labor in a hot environment. Anhidrosis may occur more readily when some disturbance of sweating already exists. If there is evidence of anhidrosis, the possibility of hyperthermia should be considered. Dosage should be decreased at the discretion of the physician so that the ability to maintain body heat equilibrium by perspiration is not impaired. Severe anhidrosis and fatal hyperthermia have occurred.

PRECAUTIONS

General

Since benztropine mesylate has cumulative action, continued supervision is advisable. Patients with a tendency to tachycardia and patients with prostatic hypertrophy should be observed closely during treatment.

Dysuria may occur, but rarely becomes a problem. Urinary retention has been reported with benztropine mesylate.

The drug may cause complaints of weakness and inability to move particular muscle groups, especially in large doses. For example, if the neck has been rigid and suddenly relaxes, it may feel weak, causing some concern. In this event, dosage adjustment is required.

Mental confusion and excitement may occur with large doses, or in susceptible patients. Visual hallucinations have been reported occasionally. Furthermore, in the treatment of extrapyramidal disorders due to neuroleptic drugs (e.g., phenothiazines), in patients with mental disorders, occasionally there may be intensification of mental symptoms. In such cases, antiparkinsonian drugs can precipitate a toxic psychosis. Patients with mental disorders should be kept under careful observation, especially at the beginning of treatment or if dosage is increased.

Tardive dyskinesia may appear in some patients on long-term therapy with phenothiazines and related agents, or may occur after therapy with these drugs has been discontinued. Antiparkinsonism agents do not alleviate the symptoms of tardive dyskinesia, and in some instances may aggravate them. Benztropine mesylate is not recommended for use in patients with tardive dyskinesia.

The physician should be aware of the possible occurrence of glaucoma. Although the drug does not appear to have any adverse effect on simple glaucoma, it probably should not be used in angle-closure glaucoma.

Drug Interactions

Antipsychotic drugs such as phenothiazines or haloperidol; tricyclic antidepressants (see **WARNINGS**).

Pediatric Use

Because of the atropine-like side effects, benztropine mesylate should be used with caution in pediatric patients over three years of age (see **CONTRAINDICATIONS**).

ADVERSE REACTIONS

The adverse reactions below, most of which are anticholinergic in nature, have been reported and within each category are listed in order of decreasing severity.

Cardiovascular

Tachycardia.

Digestive

Paralytic ileus, constipation, vomiting, nausea, dry mouth.

If dry mouth is so severe that there is difficulty in swallowing or speaking, or loss of appetite and weight, reduce dosage, or discontinue the drug temporarily.

Slight reduction in dosage may control nausea and still give sufficient relief of symptoms. Vomiting may be controlled by temporary discontinuation, followed by resumption at a lower dosage.

Nervous System

Toxic psychosis, including confusion, disorientation, memory impairment, visual hallucinations; exacerbation of preexisting psychotic symptoms; nervousness; depression; listlessness; numbness of fingers.

Special Senses

Blurred vision, dilated pupils.

Urogenital

Urinary retention, dysuria.

Metabolic/Immune or Skin

Occasionally, an allergic reaction, e.g., skin rash, develops. If this cannot be controlled by dosage reduction, the medication should be discontinued.

Other

Heat stroke, hyperthermia, fever.

OVERDOSAGE

Manifestations - May be any of those seen in atropine poisoning or antihistamine overdose: CNS depression, preceded or followed by stimulation; confusion; nervousness; listlessness; intensification of mental symptoms or toxic psychosis in patients with mental illness being treated with neuroleptic drugs (e.g., phenothiazines); hallucinations (especially visual); dizziness; muscle weakness; ataxia; dry mouth; mydriasis; blurred vision; palpitations; tachycardia; elevated blood pressure; nausea; vomiting; dysuria; numbness of fingers; dysphagia; allergic reactions, e.g., skin rash; headache; hot, dry, flushed skin; delirium; coma; shock; convulsions; respiratory arrest; anhidrosis; hyperthermia; glaucoma; constipation.

Treatment - Physostigmine salicylate, 1 to 2 mg, SC or IV, reportedly will reverse symptoms of anticholinergic intoxication.* A second injection may be given after 2 hours if required. Otherwise treatment is symptomatic and supportive. Induce emesis or perform gastric lavage (contraindicated in precomatose, convulsive, or psychotic states). Maintain respiration. A short-acting barbiturate may be used for CNS excitement, but with caution to avoid subsequent depression; supportive care for depression (avoid convulsant stimulants such as picrotoxin, pentylenetetrazol, or bemegride); artificial respiration for severe respiratory depression; a local miotic for mydriasis and cycloplegia; ice bags or other cold applications and alcohol sponges for hyperpyrexia, a vasopressor and fluids for circulatory collapse. Darken room for photophobia.

DOSAGE AND ADMINISTRATION

Benztropine mesylate tablets should be used when patients are able to take oral medication.

The injection is especially useful for psychotic patients with acute dystonic reactions or other reactions that make oral medication difficult or impossible. It is recommended also

when a more rapid response is desired than can be obtained with the tablets.

Because of cumulative action, therapy should be initiated with a low dose which is increased gradually at five or six-day intervals to the smallest amount necessary for optimal relief. Increases should be made in increments of 0.5 mg, to a maximum of 6 mg, or until optimal results are obtained without excessive adverse reactions.

Postencephalitic and Idiopathic Parkinsonism - The usual daily dose is 1 to 2 mg, with a range of 0.5 to 6 mg orally or parenterally.

As with any agent used in parkinsonism, dosage must be individualized according to age and weight, and the type of parkinsonism being treated. Generally, older patients, and thin patients cannot tolerate large doses. Most patients with postencephalitic parkinsonism need fairly large doses and tolerate them well. Patients with a poor mental outlook are usually poor candidates for therapy.

In idiopathic parkinsonism, therapy may be initiated with a single daily dose of 0.5 to 1 mg at bedtime. In some patients, this will be adequate; in others 4 to 6 mg a day may be required.

In postencephalitic parkinsonism, therapy may be initiated in most patients with 2 mg a day in one or more doses. In highly sensitive patients, therapy may be initiated with 0.5 mg at bedtime, and increased as necessary.

Some patients experience greatest relief by taking the entire dose at bedtime; others react more favorably to divided doses, two to four times a day. Frequently, one dose a day is sufficient, and divided doses may be unnecessary or undesirable.

The long duration of action of this drug makes it particularly suitable for bedtime medication when its effects may last throughout the night, enabling patients to turn in bed during the night more easily, and to rise in the morning.

When benztropine mesylate is started, do not terminate therapy with other antiparkinsonian agents abruptly. If the other agents are to be reduced or discontinued, it must be done gradually. Many patients obtain greatest relief with combination therapy.

Benztropine mesylate may be used concomitantly with Carbidopa-Levodopa, or with levodopa, in which case periodic dosage adjustment may be required in order to maintain optimum response.

Drug-Induced Extrapyrarnidal Disorders - In treating extrapyramidal disorders due to neuroleptic drugs (e.g., phenothiazines), the recommended dosage is 1 to 4 mg once or twice a day orally or parenterally. Dosage must be individualized according to the need of the patient. Some patients require more than recommended; others do not need as much.

In acute dystonic reactions, 1 to 2 mL of the injection usually relieves the condition quickly. After that, the tablets, 1 to 2 mg twice a day, usually prevent recurrence.

When extrapyramidal disorders develop soon after initiation of treatment with neuroleptic drugs (e.g., phenothiazines), they are likely to be transient. One to 2 mg of benztropine mesylate tablets two or three times a day usually provides relief within one or two days. After one or two weeks, the drug should be withdrawn to determine the continued need for it. If such disorders recur, benztropine mesylate can be reinstated.

Certain drug-induced extrapyramidal disorders that develop slowly may not respond to

benztropine mesylate.

HOW SUPPLIED

Benzotropine Mesylate Tablets, USP 0.5 mg are round, white, debossed with 'CE' on the left side of the functional score and '41' on the right side on one side and plain on the other side.

They are supplied in:

Bottles of 30	NDC 62135-141-30
Bottles of 100	NDC 62135-141-01
Bottles of 180	NDC 62135-141-18
Bottles of 540	NDC 62135-141-57
Bottles of 1000	NDC 62135-141-10

Benzotropine Mesylate Tablets, USP 1 mg are oval shaped, white, debossed with 'CE' on the left side of the functional score and '44' on the right side on one side and plain on the other side.

They are supplied in:

Bottles of 30	NDC 62135-144-30
Bottles of 100	NDC 62135-144-01
Bottles of 180	NDC 62135-144-18
Bottles of 540	NDC 62135-144-57
Bottles of 1000	NDC 62135-144-10

Benzotropine Mesylate Tablets, USP 2 mg are round, white, debossed with 'CE' on the upper side of the functional score and '42' on the lower side on one side and plain on the other side.

They are supplied in:

Bottles of 30	NDC 62135-142-30
Bottles of 100	NDC 62135-142-01
Bottles of 180	NDC 62135-142-18
Bottles of 540	NDC 62135-142-57
Bottles of 1000	NDC 62135-142-10

Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].

Dispense in a tight, light-resistant container [see USP].

* Duvoisin, R.C.; Katz, R.J; Amer. Med. Ass. 206: 1963-1965, Nov. 25, 1968.

Manufactured by: Chartwell Pharmaceuticals, LLC.

Congers, NY 10920

Manufactured for: Chartwell RX, LLC.

Congers, NY 10920

Made in USA

Rev: 08/2022

L70507

PRINCIPAL DISPLAY PANEL

NDC 62135-144-01

Benztropine

Mesylate

Tablets, USP

1 mg

Rx Only

100 Tablets

NDC 62135-144-01

**Benzotropine
Mesylate
Tablets, USP**

1 mg

Rx Only
100 Tablets

Chartwell Rx

Each tablet contains 1 mg of Benzotropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for: Chartwell RX, LLC.
Congers, NY 10920

L70502
REV. 01 10/20

GTIN 00362135144012

3 6 2 1 3 5 1 4 4 0 1 2

Made in USA

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-144-18

Benzotropine

Mesylate

Tablets, USP

1 mg

Rx Only

180 Tablets

NDC 62135-144-18

Benztropine Mesylate Tablets, USP

1 mg

Rx Only
180 Tablets

Chartwell Rx

Each tablet contains 1 mg of Benzotropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].

USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for: Chartwell RX, LLC.
Congers, NY 10920

L70872  Made in U.S.A.
GTIN 00362135144180 REV. 01 03/22



N 3 62135 14418 0

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-144-57
Benztropine Mesylate Tablets, USP
1 mg
Rx Only
540 Tablets

NDC 62135-144-57

Benzotropine Mesylate Tablets, USP

1 mg

Rx Only
540 Tablets


Chartwell Rx


Each tablet contains 1 mg of Benzotropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].

USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for: Chartwell RX, LLC.
Congers, NY 10920

L70873  Made in U.S.A.
GTIN 00362135144579 REV. 01 03/22



N 3 62135 14457 9

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-144-10
Benztropine Mesylate Tablets, USP
1 mg

Rx Only
1000 Tablets

NDC 62135-144-10
Benztropine Mesylate Tablets, USP

1 mg

Rx Only
1000 Tablets

Chartwell Rx

Each tablet contains 1 mg of Bzotropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.
Manufactured for: Chartwell RX, LLC.
Congers, NY 10920 L70503
GTIN 00362135144104 REV. 01 10/20  Made in USA


N 3 6 2 1 3 5 1 4 4 1 0 4

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-142-01
Benztropine Mesylate Tablets, USP
2 mg
Rx Only
100 Tablets

NDC 62135-142-01
Benztropine Mesylate Tablets, USP

2 mg

Rx Only
100 Tablets

Chartwell Rx

Each tablet contains 2 mg of Bzotropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.
Manufactured for: Chartwell RX, LLC.
Congers, NY 10920 L70505
GTIN 00362135142018 REV. 01 10/20  Made in USA


N 3 6 2 1 3 5 1 4 2 0 1 8

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-142-18
Benztropine Mesylate

Tablets, USP
2 mg
Rx Only
180 Tablets

NDC 62135-142-18

Benztropine Mesylate Tablets, USP

2 mg

Rx Only
180 Tablets

Chartwell Rx

Each tablet contains 2 mg of Benzotropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.
Manufactured for: Chartwell RX, LLC.
Congers, NY 10920

GTIN 00362135142186 REV. 01 03/22

L70874
Made in U.S.A.



N 3 6213514218 6

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-142-57
Benzotropine Mesylate Tablets, USP
2 mg
Rx Only
540 Tablets

NDC 62135-142-57

Benzotropine Mesylate Tablets, USP

2 mg


Rx Only
540 Tablets

Chartwell Rx

Each tablet contains 2 mg of Benzotropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.
Manufactured for: Chartwell RX, LLC.
Congers, NY 10920

GTIN 00362135142575 REV. 01 03/22

L70875
Made in U.S.A.



N 3 6213514257 5

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-142-10

Benztropine
Mesylate
Tablets, USP
2 mg
Rx Only
1000 Tablets

NDC 62135-142-10

Benztropine Mesylate Tablets, USP

2 mg

Rx Only
1000 Tablets

Chartwell Rx

Each tablet contains 2 mg of Benztropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for: Chartwell RX, LLC.
Congers, NY 10920 L70506
GTIN 00362135142100 REV. 01 10/20  Made in USA



N 3 6 2 1 3 5 1 4 2 1 0 0

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-141-18
Benztropine
Mesylate
Tablets, USP
0.5 mg
Rx Only
180 Tablets

NDC 62135-141-18

Benztropine Mesylate Tablets, USP

0.5 mg

Rx Only
180 Tablets

Chartwell Rx

Each tablet contains 0.5 mg of Benztropine Mesylate, USP.
Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature].
Dispense in a tight, light-resistant container [see USP].
USUAL ADULT DOSAGE: For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.
KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for: Chartwell RX, LLC.
Congers, NY 10920 L70870
GTIN 00362135141189 REV. 01 03/22  Made in USA



N 3 6 2 1 3 5 1 4 1 1 8 9

No Varnish

PRINCIPAL DISPLAY PANEL

NDC 62135-141-57
 Bzotropine
 Mesylate
 Tablets, USP
 0.5 mg
 Rx Only
 540 Tablets

NDC 62135-141-57
Benzotropine Mesylate
 Tablets, USP

0.5 mg

Rx Only
 540 Tablets

Chartwell Rx

Each tablet contains 0.5 mg of Benzotropine Mesylate, USP. Store at 20° to 25°C (68° to 77°F); excursions permitted to 15° to 30°C (59° to 86°F) [see USP Controlled Room Temperature]. Dispense in a tight, light-resistant container [see USP].

USUAL ADULT DOSAGE For parkinsonism, 1 to 2 mg daily. For drug induced extrapyramidal disorders, 1 to 4 mg once or twice a day. See accompanying insert.

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Manufactured for: Chartwell RX, LLC.
 Congers, NY 10920

L70871
 REV. 01 03/22
 Made in U.S.A.

GTIN 00362135141578

3 62135 14157 8

No Varnish

BENZTROPINE MESYLATE

benztropine mesylate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62135-141
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZTROPINE MESYLATE (UNII: VMJ8TL7510) (BENZTROPINE - UNII:1NHL2J4X8K)	BENZTROPINE MESYLATE	0.5 mg

Inactive Ingredients

Ingredient Name	Strength
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	7mm
Flavor		Imprint Code	CE;41
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62135-141-18	180 in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2022	
2	NDC:62135-141-57	540 in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA081265	01/23/1992	

BENZTROPINE MESYLATE

benztropine mesylate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62135-144
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZTROPINE MESYLATE (UNII: WMJ8TL7510) (BENZTROPINE - UNII:1NHL2J4X8K)	BENZTROPINE MESYLATE	1 mg

Inactive Ingredients

Ingredient Name	Strength
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	CE;44

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62135-144-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2021	
2	NDC:62135-144-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2021	
3	NDC:62135-144-18	180 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/07/2022	
4	NDC:62135-144-57	540 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/07/2022	
5	NDC:62135-144-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA081265	01/23/1992	

BENZTROPINE MESYLATE

benztropine mesylate tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:62135-142
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZTROPINE MESYLATE (UNII: WMJ8TL7510) (BENZTROPINE - UNII:1NHL2J4X8K)	BENZTROPINE MESYLATE	2 mg

Inactive Ingredients

Ingredient Name	Strength
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	white	Score	2 pieces
--------------	-------	--------------	----------

Shape	ROUND	Size	7mm
Flavor		Imprint Code	CE;42
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62135-142-30	30 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2021	
2	NDC:62135-142-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2021	
3	NDC:62135-142-18	180 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/07/2022	
4	NDC:62135-142-57	540 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/07/2022	
5	NDC:62135-142-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/30/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA081265	01/23/1992	

Labeler - Chartwell RX, LLC. (079394054)

Revised: 1/2023

Chartwell RX, LLC.