

ADULT CHEWABLE ASPIRIN- aspirin tablet, chewable HARRIS TEETER

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1152B-HTE-2021-0906

Drug Facts

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- temporarily relieves minor aches and pains
- ask your doctor about other uses for this product

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: do not use unless directed by a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicon dioxide, sucrose

Questions or comments?

1-844-705-4384

PRINCIPAL DISPLAY PANEL

Harris Teeter®

NDC 72036-252-01

Adult Chewable Aspirin

†Compare to the Active Ingredient in St. Joseph® Low Dose Chewable Aspirin

Low Dose, 81 mg

PAIN RELIEVER (NSAID)

- Orange Flavored
- Adult Aspirin Regimen**

Actual Size

36 CHEWABLE TABLETS

INK AND COATING FREE
FOR LOT AND
EXPIRATION STAMPING



DO NOT USE IF IMPRINTED SEAL
UNDER CAP IS BROKEN OR MISSING

FT 152801H1E, R0
This product is not manufactured or distributed by
Foundation Consumer Healthcare, LLC, distributor of
St. Joseph® Low Dose Chewable Aspirin.

Drug Facts (continued)

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Do not use**
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
 - for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

Ask a doctor before use if

- stomach bleeding warning applies to you
 - you have a history of stomach problems, such as heartburn
 - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 - you are taking a diuretic
 - you have asthma
- Ask a doctor or pharmacist before use if you are taking a prescription drug for:**
- gout
 - diabetes
 - arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - ringing in the ears or a loss of hearing occurs
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

- corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicon dioxide, sucrose

NDC 72036-252-01

Adult Chewable Aspirin



†Compare to the Active Ingredient in St. Joseph® Low Dose Chewable Aspirin

Low Dose, 81 mg

PAIN RELIEVER (NSAID)

- Orange Flavored
- Adult Aspirin Regimen**

Actual Size



36 CHEWABLE TABLETS

Drug Facts

Active ingredient

Aspirin 81 mg (NSAID) Pain reliever
* nonsteroidal anti-inflammatory drug

Purpose

- temporarily relieves minor aches and pains
- ask your doctor about other uses for this product

Uses

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 ■ hives
 ■ asthma (wheezing)
 ■ shock
 ■ facial swelling

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug

Drug Facts (continued)

Questions or comments?
1-844-705-4384

PRODUCE DISTRIBUTED BY:
HARRIS TEETER, LLC MATTHEWS, NC 28105
1-800-432-6111 or harrisiteeter.com
** Talk to your doctor before starting an aspirin regimen. Aspirin is not right for everyone.

ADULT CHEWABLE ASPIRIN aspirin tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72036-252
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCROSE (UNII: C151H8M554)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	AP;009
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72036-252-01	1 in 1 CARTON	09/06/2021	
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/06/2021	

Labeler - HARRIS TEETER (047279351)

Revised: 9/2021

HARRIS TEETER