

**ANTI-DIARRHEAL- loperamide hcl tablet**  
**Rite Aid Corporation**

-----  
**Rite Aid 44-375**

***Active ingredient (in each caplet)***

Loperamide HCl 2 mg

***Purpose***

Anti-diarrheal

***Use***

controls symptoms of diarrhea, including Travelers' Diarrhea

***Warnings***

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use**

if you have bloody or black stool.

**Ask a doctor before use if you have**

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product,**

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

- store between 20°-25°C (68°-77°F)
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- see end flap for expiration date and lot number

**Inactive ingredients**

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

**Questions or comments?**

Call **1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

**Principal Display Panel**

**NDC 11822-0375-3**

Compare to the active ingredient in **Imodium® A-D\***

**ANTI-DIARRHEAL**

**LOPERAMIDE HCl TABLETS, 2 mg**

ANTI-DIARRHEAL  
FOR THE CONTROL OF THE  
SYMPTOMS OF DIARRHEA

**ACTUAL SIZE**

**12  
CAPLETS**

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR  
SHOWS ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Imodium® A-D. 50844 REV0619K37502

**DISTRIBUTED BY:** RITE AID,  
200 NEWBERRY COMMONS  
ETTERS, PA 17319  
**[www.riteaid.com](http://www.riteaid.com)**

**SATISFACTION  
GUARANTEE**

If you're not satisfied, we'll  
happily refund your money.

1. To open, tear at perforations and remove a section.
2. Use ✂ to cut through plastic for caplet.

**Drug Facts**  
**Active ingredient (in each caplet)** Loperamide HCl 2 mg  
**Purpose** Anti-diarrheal  
**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

NDC 11822-0375-3

Compare to the active ingredient in **Imodium® A-D\***

# ANTI-DIARRHEAL

**LOPERAMIDE HCl TABLETS, 2 mg**

ANTI-DIARRHEAL  
 FOR THE CONTROL OF THE  
 SYMPTOMS OF DIARRHEA



ACTUAL SIZE

**12**  
 CAPLETS



**SATISFACTION GUARANTEE**  
 If you're not satisfied, we'll happily refund your money.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DISTRIBUTED BY: RITE AID  
 200 NEUBERRY COMMONS  
 ETTERS, PA 17319  
 www.riteaid.com

\*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Imodium® A-D. 50844 REV0619K37502

No print/No varnish area  
 Lot no/Exp date



1. To open, tear at perforations and remove a section.



2. Use caplet cutter to cut through plastic for caplet.

**Drug Facts (continued)**  
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
**Inactive ingredients** corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide  
**Questions or comments?** Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

B-1702-375-02-R3  
 REV0619K37502

**Other information** ■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

■ find right dose on chart. If possible, use weight to dose; otherwise, use age.  
**Directions** ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea  
 ■ Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.  
 ■ If pregnant or breast-feeding, ask a health professional before use.  
 ■ you get abdominal swelling or binging. These may be signs of a serious condition.  
**Stop use and ask a doctor if** ■ symptoms get worse ■ diarrhea lasts for more than 2 days  
 ■ When using this product, tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.  
 ■ Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

**Drug Facts (continued)****Warnings:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.**Heart alert:** Taking more than directed can cause serious heart problems or death.

Do not use if you have bloody or black stool.

Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm

**Rite Aid 44-375****ANTI-DIARRHEAL**

loperamide hcl tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-0375
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

**Product Characteristics**

<b>Color</b>	green (light)	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;375
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-0375-8	4 in 1 CARTON	05/03/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

2	NDC:11822-0375-3	2 in 1 CARTON	05/03/2005	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:11822-0375-5	1 in 1 CARTON	05/03/2005	07/15/2021
3		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/03/2005	

**Labeler** - Rite Aid Corporation (014578892)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(11822-0375) , pack(11822-0375)

## Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(11822-0375)

Revised: 4/2024

Rite Aid Corporation