

**FOSTER AND THRIVE STOMACH RELIEF - bismuth subsalicylate tablet**  
**Strategic Sourcing Services LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Foster and Thrive Stomach Relief caplets 172**

**ACTIVE INGREDIENT(in each caplet)**

Bismuth subsalicylate 262 mg

**PURPOSE**

Upset stomach reliever and anti-diarrheal

**USE(S)**

relieves:

- indigestion
- nausea
- heartburn
- diarrhea
- upset stomach associated with these symptoms

**WARNINGS**

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**DO NOT USE IF YOU HAVE**

- bloody or black stool
- an ulcer
- a bleeding problem

## **ASK A DOCTOR BEFORE USE IF YOU HAVE**

- fever
- mucus in the stool

## **ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

## **WHEN USING THIS PRODUCT**

a temporary, but harmless darkening of the stool and/or tongue may occur.

## **STOP USE AND ASK DOCTOR IF**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **IF PREGNANT OR BREAST-FEEDING,**

ask a health professional before use.

## **KEEP OUT OF REACH OF CHILDREN.**

In case of overdose, get medical help or contact a Poison Control Center immediately(1-800-222-1222).

## **DIRECTIONS**

- swallow caplet(s) with water, do not chew
- adults and children 12 years and over: 2 caplets every 1/2 to 1 hour as needed
- do not take more than 8 doses (16 caplets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

## **OTHER INFORMATION**

- **each caplet contains:**
- calcium 27 mg, salicylate 102 mg and sodium 2 mg
- very low sodium
- **TAMPER EVIDENT: do not use if printed seal under cap is torn or missing**

- store at room temperature, in a dry place

## **INACTIVE INGREDIENTS**

calcium carbonate, colloidal silicon dioxide, D&C red 27 lake, magnesium stearate, mannitol, microcrystalline cellulose, polysorbate 80, sodium starch glycolate.

## **QUESTIONS OR COMMENTS?**

**Call 833-358-6431** Monday to Friday 9:00 am to 7:00 pm EST

## **PRINCIPAL DISPLAY PANEL**

NDC 70677-1082-1

COMPATE TO PEPTO-BISMOL<sup>®</sup> CAPLETS ACTIVE INGREDIENT\*

### **Foster and Thrive TM**

#### **Stomach Relief**

**Bismuth Subsalicylate 262 mg**

**ANTIDIARRHEAL / UPSET STOMACH RELIEVER**

#### **Relieves:**

- Upset Stomach
- Nausea
- Heartburn
- Indigestion
- Diarrhea

40 CAPLETS

©2023 McKesson Corporation  
 MFR# 53041 REV 0323  
**Active ingredient** (in each caplet): Bismuth subsalicylate (262 mg)  
**Uses relieves:** ■ indigestion ■ nausea ■ heartburn ■ diarrhea ■ upset stomach  
**Warnings:** **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Contains salicylate. Do not take if you are allergic to salicylates (including aspirin) ■ taking other salicylate products. Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool.

NDC 70677-1082-1  
**Foster & Thrive™**  
**Stomach Relief**  
**Bismuth Subsalicylate**  
**262 mg**  
**40 CAPLETS**

**REFER TO CARTON FOR FULL LABELING INFORMATION**

**Ask a doctor before use if you have** ■ fever ■ mucus in the stool. **Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis. **When using this product a temporary, but harmless darkening of the stool and/or tongue may occur. Stop use and ask a doctor if** ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days. **If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.**

**Directions** ■ swallow caplets) with water; do not chew ■ adults and children 12 years and over: 2 caplets every 1/2 to 1 hour as needed ■ do not take more than 8 doses (16 caplets) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea.

**Other information each caplet contains:** calcium 27 mg, salicylate 102 mg and sodium 2 mg. ■ **TAMPER EVIDENT:** do not use if printed seal under cap is torn or missing ■ store at room temperature, in a dry place.

Distributed by: McKesson Corp., via SSSL  
 Memphis, TN 38141  
 Money Back Guarantee [www.fosterandthrive.com](http://www.fosterandthrive.com)

Blank Area for LOT & EXP. Information

LOT: \_\_\_\_\_  
 EXP: \_\_\_\_\_



**FOSTER AND THRIVE STOMACH RELIEF**  
 bismuth subsalicylate tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70677-1082
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

## Product Characteristics

<b>Color</b>	PINK	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	G172
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:70677-1082-1	1 in 1 CARTON	05/15/2023	
1		40 in 1 BOTTLE; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC MONOGRAPH FINAL	part335	05/15/2023	

Revised: 5/2023

Strategic Sourcing Services LLC