

# **STOMACH RELIEF ULTRA- bismuth subsalicylate liquid QUALITY CHOICE (Chain Drug Marketing Association)**

-----  
**Drug Facts**

**Active ingredient (in each 15 mL)**

Bismuth subsalicylate 525 mg

**Purpose**

Upset stomach reliever/Antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - gas
  - indigestion
  - nausea
  - fullness
  - belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use**

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool

## **Ask a doctor or pharmacist before use if you are**

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

## **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur.

## **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **If pregnant or breast-feeding,**

ask a health professional before use.

## **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

## **Directions**

- do not take more than 8 doses (120 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
  - adults and children 12 years and over:
    - 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
    - 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- children under 12 years of age: ask a doctor

## **Other information**

- **each 15 mL contains:** sodium 6 mg
- **each 15 mL contains:** salicylate 227 mg
- low sodium
- keep tightly closed

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

### **Inactive ingredients**

benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid

### **Questions or comments?**

Call **1-800-935-2362** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

\*Compare to the Active Ingredient IN **Pepto-Bismol® Ultra**

Ultra

### **Stomach Relief**

Stomach Relief

Bismuth subsalicylate | Upset Stomach Reliever/Antidiarrheal

For Relief of:

Heartburn

Indigestion

Nausea

Upset Stomach

Diarrhea

### **2x strength per ounce†**

Alcohol free

Sugar free

FL OZ (mL)

\*This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

**TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**

Distributed by C.D.M.A., Inc. ©

43157 W 9 Mile Rd

Novi, MI 48375

[www.qualitychoice.com](http://www.qualitychoice.com)

# Package Label

**Drug Facts** (continued)

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** ■ do not take more than 8 doses (120 mL) in 24 hours  
 ■ use until diarrhea stops but not more than 2 days ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ mL = milliliter ■ shake well before using ■ measure only with dosing cup provided. Do not use any other dosing device. ■ keep dosing cup with product ■ adults and children 12 years and over: ■ 15 mL (1 dose) every 1/2 hour or 30 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 15 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ children under 12 years of age: ask a doctor

**Other information** ■ each 15 mL contains: sodium 6 mg  
 ■ each 15 mL contains: salicylate 227 mg ■ low sodium ■ keep tightly closed ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)

**Inactive ingredients** benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid

**Questions or comments?**  
 Call 1-800-935-2362 Monday-Friday 9AM-5PM EST

Distributed by C.D.M.A., Inc.  
 43157 W 9 Mile Rd  
 Novi, MI 48375  
 www.qualitychoice.com  
 Questions: 800-935-2362

\*This product is not manufactured or distributed by The Procter & Gamble Company.  
 Pepto Bismol® is a registered trademark of The Procter & Gamble Company.  
 vs. Original Pepto-Bismol®

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.**



PLD-A559A LB008416 6 3551598132 4



NDC 63868-814-08

**"Compare to the Active Ingredient in Pepto Bismol® Ultra"**

## Ultra Stomach Relief

### Stomach Relief

Bismuth Subsalicylate | Upset Stomach Reliever, Antidiarrheal

**For Relief of:**  
 Heartburn  
 Indigestion  
 Nausea  
 Upset Stomach  
 Diarrhea

**2x strength per ounce\***

Alcohol Free  
 Sugar Free

**8 fl oz (237 mL)**

**Drug Facts**

Active ingredient (in each 15 mL)	Purpose
Bismuth subsalicylate 525 mg.....	Upset stomach reliever/Antidiarrheal

**Uses** relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including ■ heartburn ■ indigestion ■ nausea ■ gas ■ fullness ■ belching

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert: Contains salicylate. Do not take if you are** ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

**Do not use** if you have  
 ■ an ulcer ■ a bleeding problem ■ bloody or black stool

**Ask a doctor before use if you have**  
 ■ fever ■ mucus in the stool

**Ask a doctor or pharmacist before use if you are taking any drug for** ■ diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur.

**Stop use and ask a doctor if** ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.**

## QUALITY CHOICE Ultra Stomach Relief

# STOMACH RELIEF ULTRA

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:63868-814
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 7671P0Y5NH)	
<b>MAGNESIUM ALUMINUM SILICATE</b> (UNII: 6M3P64V0NC)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	

<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>SODIUM SALICYLATE</b> (UNII: MQ1H85SYP)	
<b>SORBIC ACID</b> (UNII: X045WJ989B)	
<b>METHYLCELLULOSE (1500 CPS)</b> (UNII: P0NTE48364)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-814-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/26/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/26/2021	

**Labeler** - QUALITY CHOICE (Chain Drug Marketing Association) (011920774)

Revised: 5/2024

QUALITY CHOICE (Chain Drug Marketing Association)