

BERKLEY AND JENSEN ANTI DIARRHEAL- loperamide hydrochloride tablet, film coated
BJWC

BJWC Anti Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days

- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-800-934-1204

Principal Display Panel

Compare to the Active Ingredient in Imodium® A-D

berkly jensen®

LOPERAMIDE HYDROCHLORIDE TABLETS, 2mg

ANTI-DIARRHEAL

CONTROLS THE SYMPTOMS OF DIARRHEA

ANTI-DIARRHEAL

ACTUAL SIZE

24 CAPLETS*

(*CAPSULE-SHAPED TABLETS)

100% MONEY-BACK GUARANTEE



Questions or comments?
1-800-934-1204

Drug Facts (continued)
Inactive ingredients anhydrous lactose, camouba wax, D.C. yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch



gluten free
*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D. BJ #241158 Mfg. #075798098

OPEN OTHER END
DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

CODE AREA

22462 D3 C-2

Drug Facts
Active ingredient (in each caplet) Loperamide HCl 2 mg.....Anti-diarrheal
Purpose Use controls symptoms of diarrhea including Traveler's Diarrhea

Warnings
Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl
Heart alert: Taking more than directed can cause serious heart problems or death
Do not use if you have bloody or black stool

Ask a doctor before use if you have
■ fever
■ mucus in the stool
■ a history of liver disease
■ a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if
■ symptoms get worse
■ diarrhea lasts for more than 2 days
■ you get abdominal swelling or bulging.

These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use.

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
■ find right dose on chart, if possible, use weight to dose; otherwise, use age.

adults and children 12 years and over
2 caplets after the first loose stool; 1 caplet after each subsequent loose stool, but no more than 4 caplets in 24 hours

children 9-11 years (60-95 lbs)
1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours

children 6-8 years (48-59 lbs)
1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours

children 2-5 years (34 to 47 lbs)
ask a doctor

children under 2 years (up to 33 lbs)
do not use

Other information
■ store at 20°-25°C (68°-77°F)
■ see end panel for lot number and expiration date

Compare to the Active Ingredient in Imodium® A-D**



LOPERAMIDE HYDROCHLORIDE TABLETS, 2mg
ANTI-DIARRHEAL

Compare to the Active Ingredient in Imodium® A-D**



Every Berkley Jensen product comes with our 100% Money Back Guarantee.

Distributed by:
BJ's Wholesale Club, Inc.
350 Campus Drive
Marlborough, MA 01752

CONVENIENT RE-CLOSING TAB





BERKLEY AND JENSEN ANTI DIARRHEAL

loperamide hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68391-224
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68391-224-01	2 in 1 CARTON	04/19/2003	08/31/2021
1		96 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:68391-224-62	24 in 1 CARTON	01/15/2020	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
ANDA	ANDA075232	04/19/2003		

Labeler - BJWC (159082692)

Revised: 12/2023

BJWC