

**ANTI DIARRHEAL- loperamide hcl tablet, film coated**  
**L. Perrigo Company**

-----  
**Anti-diarrheal Drug Facts**

**Active ingredient (in each caplet)**

Loperamide HCl 2 mg

**Purpose**

Anti-diarrheal

**Use**

controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

**Heart alert:** Taking more than directed can cause serious heart problems or death

**Do not use**

if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product**

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

## Other information

- store at 20° -25° C (68° -77° F)
- see end panel for lot number and expiration date

## Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

## Questions or comments?

**1-800-719-9260**

## Principal Display Panel

Compare to Imodium<sup>®</sup> A-D

active ingredient

See New Warnings

Loperamide Hydrochloride Tablets, 2 mg

anti-diarrheal

ANTI-DIARRHEAL

Controls the symptoms of diarrhea

24 CAPLETS\*

\*Capsule-Shaped Tablets

Actual Size

Loperamide Hydrochloride  
Tablets, 2 mg

**anti-diarrheal**

NDC 0113-2224-62 ITM. / ART. 1391366

Compare to  
**Imodium® A-D**  
active ingredient

See New Warnings

Loperamide Hydrochloride  
Tablets, 2 mg

**anti-diarrheal**

**ANTI-DIARRHEAL**

• Controls the symptoms of diarrhea

**24 CAPLETS\***  
\*Capsule-Shaped Tablets



Actual Size

22462 5N C1

OPEN OTHER END

**Drug Facts**

**Active ingredient (in each caplet)** Purpose  
Loperamide HCl 2 mg ..... Anti-diarrheal

**Use** controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**  
**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl  
**Heart alert:** Taking more than directed can cause serious heart problems or death  
**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**  

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**  

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging.

 These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

■ **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**  
 ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Other information**

■ store at 20°-25°C (68°-77°F)  
 ■ see end panel for lot number and expiration date

DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN

CONVENIENT RECLOSING TAB

**Drug Facts (continued)**

**Inactive ingredients** anhydrous lactose, calcium wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

**Questions or comments?**  
1-800-719-9260

Distributed By **Perrigo**<sup>®</sup>  
 Allegan, MI 49010 perrigo.com



**ANTI DIARRHEAL**

loperamide hcl tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0113-2224
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6B30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

## Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0113-2224-68	36 in 1 CARTON	06/25/2018	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0113-2224-62	24 in 1 CARTON	12/17/2019	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	06/25/2018	

**Labeler** - L. Perrigo Company (006013346)

Revised: 12/2019

L. Perrigo Company