

ACETAMINOPHEN AND IBUPROFEN (NSAID)- acetaminophen and ibuprofen (nsaid) tablet
RETAIL BUSINESS SERVICES,LLC

Dual Action Pain Reliever
Acetaminophen and Ibuprofen (NSAID) Tablets 250mg/125mg
Contains 2 Medicines

Active Ingredients (in each caplet)

Acetaminophen 250mg

Ibuprofen 125mg (NSAID)

Purposes

Pain reliever

Pain reliever

Uses

- temporarily relieves minor aches and pains due to:
- headache ■ toothache ■ backache ■ menstrual cramps
- muscular aches ■ minor pain of arthritis

Acetaminophen liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 6 caplets in 24 hours, which is the maximum daily amount for this product
- 3 or more alcoholic drinks every day while using this product

Acetaminophen allergy alert

may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert

ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening
- rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away

NSAID stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- Take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed

adults and children 12 years and over ■ **take 2 caplets every 8 hours** while symptoms persist

children under 12 years ■ ask a doctor

■ do not take more than 6 caplets in 24 hours, unless directed by a doctor

Other information

- read all warnings and directions before use. Keep carton.
- store at 20 to 25°C (68 to 77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

carnauba wax, colloidal silicon dioxide, croscarmellose sodium, crospovidone, ferric oxide red, ferric oxide yellow, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, povidone, pregelatinized starch, sodium

lauryl sulfate, stearic acid and titanium dioxide

Questions or comments?

call **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST

PDP

CAREONE
Compare to the active ingredients
of Advil[®] Dual Action*

NSC 724919/2
250 mg/75 mg
Contains 2 Medicines
72 TABLETS
(Capsule-Shaped Tablets)

DUAL ACTION PAIN RELIEVER

Actual Size

CAREONE
Compare to the active ingredients
of Advil[®] Dual Action*

NSC 724919/2
250 mg/75 mg
Contains 2 Medicines
72 TABLETS
(Capsule-Shaped Tablets)

DUAL ACTION PAIN RELIEVER

Actual Size

CAREONE
Compare to the active ingredients
of Advil[®] Dual Action*

NSC 724919/2
250 mg/75 mg
Contains 2 Medicines
72 TABLETS
(Capsule-Shaped Tablets)

DUAL ACTION PAIN RELIEVER

Actual Size

Drug Facts (continued)

Keep out of reach of children. See important information about this medicine on the label and package.

Other information

- Keep this medicine dry and away from heat and moisture.
- Do not use if the seal is broken or missing.
- Do not use if the medicine is past its expiration date.
- Do not use if the medicine has become discolored or has a strong, unpleasant odor.
- Do not use if the medicine has become hard, soft, or sticky.
- Do not use if the medicine has become lumpy or clumpy.
- Do not use if the medicine has become chunky or crumbly.
- Do not use if the medicine has become watery or milky.
- Do not use if the medicine has become foamy or bubbly.
- Do not use if the medicine has become white or grey.
- Do not use if the medicine has become yellow or brown.
- Do not use if the medicine has become black or dark brown.
- Do not use if the medicine has become purple or blue.
- Do not use if the medicine has become pink or red.
- Do not use if the medicine has become green or black.
- Do not use if the medicine has become any other color.

Questions or comments?
Call 1-877-770-3183 Mon-Fri 8:00 AM EST to 5:00 PM PST

200000004903
700000003266

200000004903
700000003266

3 4152000630 **2**

Lot: _____
Exp.: _____



ACETAMINOPHEN AND IBUPROFEN (NSAID)

acetaminophen and ibuprofen (nsaid) tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72476-131
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	125 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg

Inactive Ingredients

Ingredient Name	Strength
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3S)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
CROSPVIDONE (UNII: 2S7830E561)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	yellow (Light yellow to yellow colored)	Score	no score
Shape	CAPSULE (capsule shaped, biconvex, film coated tablets)	Size	14mm
Flavor		Imprint Code	G;131
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72476-131-81	18 in 1 BOTTLE; Type 0: Not a Combination Product	07/20/2023	
2	NDC:72476-131-72	72 in 1 BOTTLE; Type 0: Not a Combination Product	07/20/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA216592	07/20/2023	

Labeler - RETAIL BUSINESS SERVICES,LLC (967989935)

Revised: 12/2023

RETAIL BUSINESS SERVICES,LLC