

**MOTRIN DUAL ACTION WITH TYLENOL- acetaminophen and ibuprofen tablet,
film coated**

Johnson & Johnson Consumer Inc.

Motrin® Dual Action With Tylenol®

Drug Facts

Active ingredients (in each tablet)

Acetaminophen 250 mg

Ibuprofen 125 mg (NSAID *)

Purposes

Pain reliever

Pain reliever

Uses

■ temporarily relieves minor aches and pains due to:

■ headache

■ toothache

■ backache

■ menstrual cramps

■ muscular aches

■ minor pain of arthritis

Warnings

Liver damage warning:

This product contains acetaminophen. Severe liver damage may occur if you take:

■ with other drugs containing acetaminophen

■ more than 6 tablets in 24 hours, which is the maximum daily amount for this product

■ 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen allergy alert: may cause severe skin reactions.

Symptoms may include:

■ skin reddening ■ blisters ■ rash

If skin reaction occurs, stop use and seek medical help right away.

NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives ■ facial swelling ■ asthma (wheezing)
- shock ■ skin reddening ■ rash ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to acetaminophen or any other pain reliever
- right before or after heart surgery

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

■ you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ **do not take more than directed**

adults and children 12 years and over	■ take 2 tablets every 8 hours while symptoms persist
children under 12 years	■ ask a doctor

■ do not take more than 6 tablets in 24 hours, unless directed by a doctor

Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized

starch, stearic acid, sucralose, talc, titanium dioxide

Questions?

Call **1-877-895-3665** (toll-free) or **215-273-8755** (collect) or visit www.motrin.com

PRINCIPAL DISPLAY PANEL

NDC 50580-208-12

NEW

Motrin

DUAL ACTION WITH TYLENOL

**Acetaminophen 250 mg and
Ibuprofen (NSAID) 125 mg Tablets
Pain Reliever**

actual size

120 Tablets



Do not use with other medicines containing ACETAMINOPHEN; can cause liver damage.

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts (continued)

Active ingredients (in each tablet)

Acetaminophen 250 mg (NSAID)
Ibuprofen 125 mg (NSAID)
Pain reliever
Pain reliever

Purposes

take a blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 2 or more alcoholic drinks every day while using this product
take more for a longer time than directed

Uses

temporarily relieves minor aches and pains due to
headache toothache backache menstrual cramps
muscle aches minor pain of arthritis

Warnings

This product contains acetaminophen. Severe liver damage may occur if you take
more than 6 tablets in 24 hours, which is the maximum daily amount
with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product
Acetaminophen allergy alert: may cause severe skin reactions.
Symptoms may include:
skin redness hives rash
if skin reaction occurs, stop use and seek medical help right away.
NSAID allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
Symptoms may include:
hives facial swelling asthma (wheezing)
stomach pain trouble breathing
stomach pain skin redness rash hives
if an allergic reaction occurs, stop use and seek medical help right away.
NSAID stomach bleeding warning:
This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems

Other information

read all warnings and directions before use. Keep carton.
store at 20°-25°C (68°-77°F)
avoid excessive heat above 40°C (104°F)

Drug Facts (continued)

Directions do not take more than directed

adults and children take 2 tablets every 8 hours while symptoms persist
children under 12 years ask a doctor

Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
if you have ever had an allergic reaction to acetaminophen or any other pain reliever
right before or after heart surgery
Ask a doctor before use if:
you have liver disease
stomach bleeding warning applies to you
you have problems or serious side effects from taking pain relievers
you have a history of stomach problems, such as heartburn, disease, asthma, or had a stroke
you are taking a diuretic
Ask a doctor or pharmacist before use if you are:
under a doctor's care for any serious condition
taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
taking any other drug
When using this product:
take with food or milk if stomach upset occurs
Stop use and ask a doctor if:
you experience any of the following signs of stomach bleeding:
other pain reliever

Drug Facts (continued)

Warnings

Severe stomach pain that does not get better
have stomach pain
you have symptoms of heart problems or stroke:
chest pain trouble breathing
stomach pain weakness in one part or side of body
stirred speech leg swelling
pain gets worse or lasts more than 10 days
redness or swelling is present in the painful area
any new symptoms appear

Drug Facts (continued)

Directions do not take more than directed

adults and children take 2 tablets every 8 hours while symptoms persist
children under 12 years ask a doctor

Other information

read all warnings and directions before use. Keep carton.
store at 20°-25°C (68°-77°F)
avoid excessive heat above 40°C (104°F)

Drug Facts (continued)

Directions do not take more than directed

adults and children take 2 tablets every 8 hours while symptoms persist
children under 12 years ask a doctor

Other information

read all warnings and directions before use. Keep carton.
store at 20°-25°C (68°-77°F)
avoid excessive heat above 40°C (104°F)

MOTRIN DUAL ACTION WITH TYLENOL			
acetaminophen and ibuprofen tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-208
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	125 mg
	ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
Inactive Ingredients			
	Ingredient Name		Strength
	POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)		
	MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
	CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)		
	TALC (UNII: 7SEV7J4R1U)		
	POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)		
	TITANIUM DIOXIDE (UNII: 15FX9V2JP)		
	POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)		
	CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)		
	SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		

STARCH, CORN (UNII: O8232NY3SJ)

SUCRALOSE (UNII: 96K6UQ3ZD4)

STEARIC ACID (UNII: 4ELV7Z65AP)

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	14mm
Flavor		Imprint Code	M;T
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-208-02	1 in 1 CARTON	03/20/2023	
1		20 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:50580-208-08	1 in 1 CARTON	03/20/2023	
2		80 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:50580-208-12	1 in 1 CARTON	03/20/2023	
3		120 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:50580-208-16	2 in 1 CARTON	04/01/2024	
4		1 in 1 CARTON		
4		80 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA214836	03/20/2023	

Labeler - Johnson & Johnson Consumer Inc. (878046358)

Revised: 3/2024

Johnson & Johnson Consumer Inc.