

**LIL DRUG STORE ANTI-DIARRHEAL- loperamide hydrochloride tablet**  
**Lil' Drug Store Products, Inc.**

-----  
**Anti-Diarrheal, Lil' Drug Store** ®

***Drug Facts***

**Active ingredient (in each caplet)**

Loperamide HCl, USP 2 mg

**Purpose**

Anti-diarrheal

**Use**

Controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert**

Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.

**Heart alert**

Taking more than directed can cause serious heart problems or death.

**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise use age

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9 - 11 years (60 - 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 - 8 years (48 - 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2 - 5 years (34 - 47 lbs)	ask a doctor
children under 2 (up to 33 lbs)	do not use

### Other information

- store between 68-77°F (20-25°C)

### Inactive ingredients

anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow no. 10 aluminum lake, FD&C Blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

### Questions or comments?

call toll-free **1-877-507-6516 (M-F 8AM - 4:30PM CST)**

## **PRINCIPAL DISPLAY PANEL - 2 mg Caplet Pouch Carton**

QUALITY

★ GUARANTEED ★

Compare to the Active  
Ingredient in Imodium® A-D\*

Anti-  
Diarrheal

on  
the  
Go  
PACKS

See New Warning & Directions

Anti-Diarrheal

Loperamide HCl Tablets, USP 2 mg

- Controls the Symptoms of Diarrhea

4  
Caplets\*\*  
2 POUCHES, 2 CAPLETS PER POUCH

Lil'  
Drug Store®

Compare to the Active Ingredient in Imodium® A-D

**Anti-Diarrheal**

on the **Go** PACKS

See New Warning & Directions

# Anti-Diarrheal

Loperamide HCl Tablets, USP 2 mg

• Controls the Symptoms of Diarrhea

**4** Caplets\*\*

2 POUCHES, 2 CAPLETS PER POUCH

**Lil' Drug Store**

4 Caplets\*\*

**Anti-Diarrheal**

**DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!**

**Drug Facts**

Active ingredient (in each caplet)	Purpose
Loperamide HCl, USP 2 mg	Anti-diarrheal

**Use** Controls symptoms of diarrhea, including Travelers' Diarrhea

**Warnings**

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to Loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use** if you have bloody or black stool

**Ask a doctor before use if you have**

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Loperamide may interact with certain prescription drugs.

**When using this product** tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

**LIFT PANEL FOR CONTINUED DRUG FACTS**

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

**Directions**

- drink plenty of clear fluids to help prevent dehydration caused by diarrhea
- find right dose on chart. If possible, use weight to dose; otherwise use age

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9 - 11 years (60 - 95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6 - 8 years (48 - 59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2 - 5 years (34 - 47 lbs)	ask a doctor
children under 2 (up to 33 lbs)	do not use

4 Caplets\*\*

**Anti-Diarrheal**

**Anti-Diarrheal**

**Drug Facts (continued)**

**Other information**

- store between 68-77°F (20-25°C)

**Inactive ingredients**

anhydrous lactose, croscarmellose sodium, crospovidone, D&C Yellow no. 10 aluminum lake, FD&C Blue no. 1 aluminum lake, hydrogenated vegetable oil, magnesium stearate, powdered cellulose, pregelatinized starch

**Questions or comments?**

call toll-free 1-877-507-6516 (M-F 8AM - 4:30PM CST)

**Product manufactured for:** Lil' Drug Store Products, Inc.  
9300 Earhart Lane SW, Cedar Rapids, IA 52404  
www.lildrugstore.com 97592C-US-05-19

\*This product is not manufactured or distributed by McNeil Consumer Healthcare. Lil' Drug Store Products, Inc. does not own the Imodium® trademark.

Each caplet (\*\*capsule-shaped tablet) contains Loperamide HCl, USP 2 mg

**EXP**

**LOT**

8 26529291516671993

**4 Caplets\*\***

# LIL DRUG STORE ANTI-DIARRHEAL

loperamide hydrochloride tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-9859
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>CROSPVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POWDERED CELLULOSE</b> (UNII: SMD1X3XO9M)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>GLYCERYL TRISTEARATE</b> (UNII: P6OCJ2551R)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	123
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-9859-2	1 in 1 CARTON	09/01/2014	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:66715-9859-7	50 in 1 CARTON	06/20/2016	09/12/2021
2		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074091	09/14/2012	

**Labeler** - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2022

Lil' Drug Store Products, Inc.